



$$J = 1$$

NODE=S044

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=S044203

Z MASS

NODE=S044205

NODE=S044205

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06). The fit is performed using the Z mass and width, the Z hadronic pole cross section, the ratios of hadronic to leptonic partial widths, and the Z pole forward-backward lepton asymmetries. This set is believed to be most free of correlations.

The Z-boson mass listed here corresponds to the mass parameter in a Breit-Wigner distribution with mass dependent width. The value is 34 MeV greater than the real part of the position of the pole (in the energy-squared plane) in the Z-boson propagator. Also the LEP experiments have generally assumed a fixed value of the $\gamma - Z$ interferences term based on the standard model. Keeping this term as free parameter leads to a somewhat larger error on the fitted Z mass. See ACCIARRI 00Q and ABBIENDI 04G for a detailed investigation of both these issues.

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
91.1876 ± 0.0021 OUR FIT				
91.1852 ± 0.0030	4.57M	¹ ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
91.1863 ± 0.0028	4.08M	² ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
91.1898 ± 0.0031	3.96M	³ ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
91.1885 ± 0.0031	4.57M	⁴ BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
91.1872 ± 0.0033		⁵ ABBIENDI	04G OPAL	$E_{cm}^{ee} = \text{LEP1} + 130-209$ GeV
91.272 ± 0.032 ± 0.033		⁶ ACHARD	04C L3	$E_{cm}^{ee} = 183-209$ GeV
91.1875 ± 0.0039	3.97M	⁷ ACCIARRI	00Q L3	$E_{cm}^{ee} = \text{LEP1} + 130-189$ GeV
91.151 ± 0.008		⁸ MIYABAYASHI	95 TOPZ	$E_{cm}^{ee} = 57.8$ GeV
91.74 ± 0.28 ± 0.93	156	⁹ ALITTI	92B UA2	$E_{cm}^{pp} = 630$ GeV
90.9 ± 0.3 ± 0.2	188	¹⁰ ABE	89C CDF	$E_{cm}^{pp} = 1.8$ TeV
91.14 ± 0.12	480	¹¹ ABRAMS	89B MRK2	$E_{cm}^{ee} = 89-93$ GeV
93.1 ± 1.0 ± 3.0	24	¹² ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630$ GeV

NODE=S044M

¹ ABBIENDI 01A error includes approximately 2.3 MeV due to statistics and 1.8 MeV due to LEP energy uncertainty.

² The error includes 1.6 MeV due to LEP energy uncertainty.

³ The error includes 1.8 MeV due to LEP energy uncertainty.

⁴ BARATE 00C error includes approximately 2.4 MeV due to statistics, 0.2 MeV due to experimental systematics, and 1.7 MeV due to LEP energy uncertainty.

⁵ ABBIENDI 04G obtain this result using the S-matrix formalism for a combined fit to their cross section and asymmetry data at the Z peak and their data at 130–209 GeV. The authors have corrected the measurement for the 34 MeV shift with respect to the Breit-Wigner fits.

⁶ ACHARD 04C select $e^+e^- \rightarrow Z\gamma$ events with hard initial-state radiation. Z decays to $q\bar{q}$ and muon pairs are considered. The fit results obtained in the two samples are found consistent to each other and combined considering the uncertainty due to ISR modelling as fully correlated.

⁷ ACCIARRI 00Q interpret the s-dependence of the cross sections and lepton forward-backward asymmetries in the framework of the S-matrix formalism. They fit to their cross section and asymmetry data at high energies, using the results of S-matrix fits to Z-peak data (ACCIARRI 00C) as constraints. The 130–189 GeV data constrains the γ/Z interference term. The authors have corrected the measurement for the 34.1 MeV shift with respect to the Breit-Wigner fits. The error contains a contribution of ± 2.3 MeV due to the uncertainty on the γZ interference.

⁸ MIYABAYASHI 95 combine their low energy total hadronic cross-section measurement with the ACTON 93D data and perform a fit using an S-matrix formalism. As expected, this result is below the mass values obtained with the standard Breit-Wigner parametrization.

⁹ Enters fit through W/Z mass ratio given in the W Particle Listings. The ALITTI 92B systematic error (± 0.93) has two contributions: one (± 0.92) cancels in m_W/m_Z and one (± 0.12) is noncancelling. These were added in quadrature.

¹⁰ First error of ABE 89 is combination of statistical and systematic contributions; second is mass scale uncertainty.

NODE=S044M;LINKAGE=DB

NODE=S044M;LINKAGE=AC

NODE=S044M;LINKAGE=AB

NODE=S044M;LINKAGE=KA

NODE=S044M;LINKAGE=AI

NODE=S044M;LINKAGE=AH

NODE=S044M;LINKAGE=NN

NODE=S044M;LINKAGE=M

NODE=S044M;LINKAGE=AG

NODE=S044M;LINKAGE=E

¹¹ ABRAMS 89B uncertainty includes 35 MeV due to the absolute energy measurement.

¹² ALBAJAR 89 result is from a total sample of 33 $Z \rightarrow e^+e^-$ events.

NODE=S044M;LINKAGE=I
NODE=S044M;LINKAGE=F

Z WIDTH

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note "The Z boson" and ref. LEP-SLC 06).

NODE=S044W

NODE=S044W

NODE=S044W

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.4952±0.0023 OUR FIT				
2.4948±0.0041	4.57M	¹ ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
2.4876±0.0041	4.08M	² ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
2.5024±0.0042	3.96M	³ ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
2.4951±0.0043	4.57M	⁴ BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
2.4943±0.0041		⁵ ABBIENDI	04G OPAL	$E_{cm}^{ee} = \text{LEP1} + 130-209$ GeV
2.5025±0.0041	3.97M	⁶ ACCIARRI	00Q L3	$E_{cm}^{ee} = \text{LEP1} + 130-189$ GeV
2.50 ±0.21 ±0.06		⁷ ABREU	96R DLPH	$E_{cm}^{ee} = 91.2$ GeV
3.8 ±0.8 ±1.0	188	ABE	89C CDF	$E_{cm}^{p\bar{p}} = 1.8$ TeV
2.42 $^{+0.45}_{-0.35}$	480	⁸ ABRAMS	89B MRK2	$E_{cm}^{ee} = 89-93$ GeV
2.7 $^{+1.2}_{-1.0}$ ±1.3	24	⁹ ALBAJAR	89 UA1	$E_{cm}^{p\bar{p}} = 546,630$ GeV
2.7 ±2.0 ±1.0	25	¹⁰ ANSARI	87 UA2	$E_{cm}^{p\bar{p}} = 546,630$ GeV

OCCUR=2

¹ ABBIENDI 01A error includes approximately 3.6 MeV due to statistics, 1 MeV due to event selection systematics, and 1.3 MeV due to LEP energy uncertainty.

² The error includes 1.2 MeV due to LEP energy uncertainty.

³ The error includes 1.3 MeV due to LEP energy uncertainty.

⁴ BARATE 00C error includes approximately 3.8 MeV due to statistics, 0.9 MeV due to experimental systematics, and 1.3 MeV due to LEP energy uncertainty.

⁵ ABBIENDI 04G obtain this result using the S-matrix formalism for a combined fit to their cross section and asymmetry data at the Z peak and their data at 130–209 GeV. The authors have corrected the measurement for the 1 MeV shift with respect to the Breit-Wigner fits.

⁶ ACCIARRI 00Q interpret the s-dependence of the cross sections and lepton forward-backward asymmetries in the framework of the S-matrix formalism. They fit to their cross section and asymmetry data at high energies, using the results of S-matrix fits to Z-peak data (ACCIARRI 00C) as constraints. The 130–189 GeV data constrains the γ/Z interference term. The authors have corrected the measurement for the 0.9 MeV shift with respect to the Breit-Wigner fits.

⁷ ABREU 96R obtain this value from a study of the interference between initial and final state radiation in the process $e^+e^- \rightarrow Z \rightarrow \mu^+\mu^-$.

⁸ ABRAMS 89B uncertainty includes 50 MeV due to the miniSAM background subtraction error.

⁹ ALBAJAR 89 result is from a total sample of 33 $Z \rightarrow e^+e^-$ events.

¹⁰ Quoted values of ANSARI 87 are from direct fit. Ratio of Z and W production gives either $\Gamma(Z) < (1.09 \pm 0.07) \times \Gamma(W)$, CL = 90% or $\Gamma(Z) = (0.82^{+0.19}_{-0.14} \pm 0.06) \times \Gamma(W)$. Assuming Standard-Model value $\Gamma(W) = 2.65$ GeV then gives $\Gamma(Z) < 2.89 \pm 0.19$ or $= 2.17^{+0.50}_{-0.37} \pm 0.16$.

NODE=S044W;LINKAGE=DB

NODE=S044W;LINKAGE=AC
NODE=S044W;LINKAGE=AB
NODE=S044W;LINKAGE=KA

NODE=S044W;LINKAGE=AI

NODE=S044W;LINKAGE=NN

NODE=S044W;LINKAGE=M

NODE=S044W;LINKAGE=I

NODE=S044W;LINKAGE=F
NODE=S044W;LINKAGE=E

Z DECAY MODES

NODE=S044215;NODE=S044

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 e^+e^-	(3.363 ±0.004) %	DESIG=1
Γ_2 $\mu^+\mu^-$	(3.366 ±0.007) %	DESIG=2
Γ_3 $\tau^+\tau^-$	(3.370 ±0.008) %	DESIG=8
Γ_4 $\ell^+\ell^-$	[a] (3.3658±0.0023) %	DESIG=16
Γ_5 $\ell^+\ell^- \ell^+\ell^-$	[b] (4.2 $^{+0.9}_{-0.8}$) × 10 ⁻⁶	DESIG=82
Γ_6 invisible	(20.00 ±0.06) %	DESIG=9
Γ_7 hadrons	(69.91 ±0.06) %	DESIG=7
Γ_8 $(u\bar{u} + c\bar{c})/2$	(11.6 ±0.6) %	DESIG=21
Γ_9 $(d\bar{d} + s\bar{s} + b\bar{b})/3$	(15.6 ±0.4) %	DESIG=22
Γ_{10} $c\bar{c}$	(12.03 ±0.21) %	DESIG=17
Γ_{11} $b\bar{b}$	(15.12 ±0.05) %	DESIG=6

Γ ₁₂	$b\bar{b}b\bar{b}$	(3.6 ± 1.3) × 10 ⁻⁴		DESIG=73
Γ ₁₃	ggg	< 1.1	% CL=95%	DESIG=64
Γ ₁₄	$\pi^0\gamma$	< 5.2	× 10 ⁻⁵ CL=95%	DESIG=10
Γ ₁₅	$\eta\gamma$	< 5.1	× 10 ⁻⁵ CL=95%	DESIG=11
Γ ₁₆	$\omega\gamma$	< 6.5	× 10 ⁻⁴ CL=95%	DESIG=48
Γ ₁₇	$\eta'(958)\gamma$	< 4.2	× 10 ⁻⁵ CL=95%	DESIG=12
Γ ₁₈	$\gamma\gamma$	< 5.2	× 10 ⁻⁵ CL=95%	DESIG=13
Γ ₁₉	$\gamma\gamma\gamma$	< 1.0	× 10 ⁻⁵ CL=95%	DESIG=14
Γ ₂₀	$\pi^\pm W^\mp$	[c] < 7	× 10 ⁻⁵ CL=95%	DESIG=18
Γ ₂₁	$\rho^\pm W^\mp$	[c] < 8.3	× 10 ⁻⁵ CL=95%	DESIG=19
Γ ₂₂	$J/\psi(1S)X$	(3.51 ^{+0.23} _{-0.25}) × 10 ⁻³	S=1.1	DESIG=23
Γ ₂₃	$\psi(2S)X$	(1.60 ± 0.29) × 10 ⁻³		DESIG=60
Γ ₂₄	$\chi_{c1}(1P)X$	(2.9 ± 0.7) × 10 ⁻³		DESIG=42
Γ ₂₅	$\chi_{c2}(1P)X$	< 3.2	× 10 ⁻³ CL=90%	DESIG=65
Γ ₂₆	$\Upsilon(1S)X + \Upsilon(2S)X$ $+ \Upsilon(3S)X$	(1.0 ± 0.5) × 10 ⁻⁴		DESIG=69
Γ ₂₇	$\Upsilon(1S)X$	< 4.4	× 10 ⁻⁵ CL=95%	DESIG=66
Γ ₂₈	$\Upsilon(2S)X$	< 1.39	× 10 ⁻⁴ CL=95%	DESIG=67
Γ ₂₉	$\Upsilon(3S)X$	< 9.4	× 10 ⁻⁵ CL=95%	DESIG=68
Γ ₃₀	$(D^0/\bar{D}^0)X$	(20.7 ± 2.0) %		DESIG=43
Γ ₃₁	$D^\pm X$	(12.2 ± 1.7) %		DESIG=44
Γ ₃₂	$D^{*}(2010)^\pm X$	[c] (11.4 ± 1.3) %		DESIG=24
Γ ₃₃	$D_{s1}(2536)^\pm X$	(3.6 ± 0.8) × 10 ⁻³		DESIG=75
Γ ₃₄	$D_{sJ}(2573)^\pm X$	(5.8 ± 2.2) × 10 ⁻³		DESIG=76
Γ ₃₅	$D^{*J}(2629)^\pm X$	searched for		DESIG=74;OUR EVAL;→ UNCHECKED ←
Γ ₃₆	BX			DESIG=61
Γ ₃₇	B^*X			DESIG=62
Γ ₃₈	B^+X	[d] (6.08 ± 0.13) %		DESIG=77
Γ ₃₉	B_s^0X	[d] (1.59 ± 0.13) %		DESIG=49
Γ ₄₀	$B_c^\pm X$	searched for		DESIG=70;OUR EVAL;→ UNCHECKED ←
Γ ₄₁	$\Lambda_c^+ X$	(1.54 ± 0.33) %		DESIG=78
Γ ₄₂	$\Xi_c^0 X$	seen		DESIG=80;OUR EST;→ UNCHECKED ←
Γ ₄₃	$\Xi_b^- X$	seen		DESIG=81;OUR EST;→ UNCHECKED ←
Γ ₄₄	b -baryon X	[d] (1.38 ± 0.22) %		DESIG=79
Γ ₄₅	anomalous γ + hadrons	[e] < 3.2	× 10 ⁻³ CL=95%	DESIG=31
Γ ₄₆	$e^+e^-\gamma$	[e] < 5.2	× 10 ⁻⁴ CL=95%	DESIG=3
Γ ₄₇	$\mu^+\mu^-\gamma$	[e] < 5.6	× 10 ⁻⁴ CL=95%	DESIG=4
Γ ₄₈	$\tau^+\tau^-\gamma$	[e] < 7.3	× 10 ⁻⁴ CL=95%	DESIG=29
Γ ₄₉	$\ell^+\ell^-\gamma\gamma$	[f] < 6.8	× 10 ⁻⁶ CL=95%	DESIG=45
Γ ₅₀	$q\bar{q}\gamma\gamma$	[f] < 5.5	× 10 ⁻⁶ CL=95%	DESIG=46
Γ ₅₁	$\nu\bar{\nu}\gamma\gamma$	[f] < 3.1	× 10 ⁻⁶ CL=95%	DESIG=47
Γ ₅₂	$e^\pm\mu^\mp$	LF [c] < 1.7	× 10 ⁻⁶ CL=95%	DESIG=5
Γ ₅₃	$e^\pm\tau^\mp$	LF [c] < 9.8	× 10 ⁻⁶ CL=95%	DESIG=25
Γ ₅₄	$\mu^\pm\tau^\mp$	LF [c] < 1.2	× 10 ⁻⁵ CL=95%	DESIG=26
Γ ₅₅	pe	L,B < 1.8	× 10 ⁻⁶ CL=95%	DESIG=71
Γ ₅₆	$p\mu$	L,B < 1.8	× 10 ⁻⁶ CL=95%	DESIG=72

[a] ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

LINKAGE=DXX

[b] Here ℓ indicates e or μ .

LINKAGE=LEM

[c] The value is for the sum of the charge states or particle/antiparticle states indicated.

LINKAGE=SG

[d] This value is updated using the product of (i) the $Z \rightarrow b\bar{b}$ fraction from this listing and (ii) the b -hadron fraction in an unbiased sample of weakly decaying b -hadrons produced in Z -decays provided by the Heavy Flavor Averaging Group (HFAG, http://www.slac.stanford.edu/xorg/hfag/osc/PDG_2009/#FRACZ).

LINKAGE=HFF

[e] See the Particle Listings below for the γ energy range used in this measurement.

LINKAGE=GDZ

[f] For $m_{\gamma\gamma} = (60 \pm 5)$ GeV.

LINKAGE=DYY

Z PARTIAL WIDTHS **$\Gamma(e^+e^-)$** **Γ_1**

For the LEP experiments, this parameter is not directly used in the overall fit but is derived using the fit results; see the note "The Z boson" and ref. LEP-SLC 06.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
83.91±0.12 OUR FIT				
83.66±0.20	137.0K	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
83.54±0.27	117.8k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
84.16±0.22	124.4k	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
83.88±0.19		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
82.89±1.20±0.89		¹ ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV

¹ ABE 95J obtain this measurement from Bhabha events in a restricted fiducial region to improve systematics. They use the values 91.187 and 2.489 GeV for the Z mass and total decay width to extract this partial width.

 $\Gamma(\mu^+\mu^-)$ **Γ_2**

This parameter is not directly used in the overall fit but is derived using the fit results; see the note "The Z boson" and ref. LEP-SLC 06.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
83.99±0.18 OUR FIT				
84.03±0.30	182.8K	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
84.48±0.40	157.6k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
83.95±0.44	113.4k	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
84.02±0.28		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

 $\Gamma(\tau^+\tau^-)$ **Γ_3**

This parameter is not directly used in the overall fit but is derived using the fit results; see the note "The Z boson" and ref. LEP-SLC 06.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
84.08±0.22 OUR FIT				
83.94±0.41	151.5K	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
83.71±0.58	104.0k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
84.23±0.58	103.0k	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
84.38±0.31		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

 $\Gamma(\ell^+\ell^-)$ **Γ_4**

In our fit $\Gamma(\ell^+\ell^-)$ is defined as the partial Z width for the decay into a pair of massless charged leptons. This parameter is not directly used in the 5-parameter fit assuming lepton universality but is derived using the fit results. See the note "The Z boson" and ref. LEP-SLC 06.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
83.984±0.086 OUR FIT				
83.82 ±0.15	471.3K	ABBIENDI	01A OPAL	$E_{cm}^{ee} = 88-94$ GeV
83.85 ±0.17	379.4k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
84.14 ±0.17	340.8k	ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
84.02 ±0.15	500k	BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

 $\Gamma(\text{invisible})$ **Γ_6**

We use only direct measurements of the invisible partial width using the single photon channel to obtain the average value quoted below. OUR FIT value is obtained as a difference between the total and the observed partial widths assuming lepton universality.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
499.0± 1.5 OUR FIT				
503 ±16 OUR AVERAGE Error includes scale factor of 1.2.				
498 ±12 ±12	1791	ACCIARRI	98G L3	$E_{cm}^{ee} = 88-94$ GeV
539 ±26 ±17	410	AKERS	95C OPAL	$E_{cm}^{ee} = 88-94$ GeV
450 ±34 ±34	258	BUSKULIC	93L ALEP	$E_{cm}^{ee} = 88-94$ GeV
540 ±80 ±40	52	ADEVA	92 L3	$E_{cm}^{ee} = 88-94$ GeV

NODE=S044218

NODE=S044W1

NODE=S044W1

NODE=S044W1

NODE=S044W1;LINKAGE=KG

NODE=S044W2

NODE=S044W2

NODE=S044W2

NODE=S044W3

NODE=S044W3

NODE=S044W3

NODE=S044W4

NODE=S044W4

NODE=S044W4

NODE=S044W6

NODE=S044W6

NODE=S044W6

• • • We do not use the following data for averages, fits, limits, etc. • • •

498.1 ± 2.6	¹ ABBIENDI	01A	OPAL	$E_{cm}^{ee} = 88-94$ GeV
498.1 ± 3.2	¹ ABREU	00F	DLPH	$E_{cm}^{ee} = 88-94$ GeV
499.1 ± 2.9	¹ ACCIARRI	00C	L3	$E_{cm}^{ee} = 88-94$ GeV
499.1 ± 2.5	¹ BARATE	00C	ALEP	$E_{cm}^{ee} = 88-94$ GeV

¹ This is an indirect determination of $\Gamma(\text{invisible})$ from a fit to the visible Z decay modes.

NODE=S044W6;LINKAGE=B1

$\Gamma(\text{hadrons})$

Γ_7

This parameter is not directly used in the 5-parameter fit assuming lepton universality, but is derived using the fit results. See the note "The Z boson" and ref. LEP-SLC 06.

NODE=S044W5

NODE=S044W5

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1744.4 ± 2.0 OUR FIT				
1745.4 ± 3.5	4.10M	ABBIENDI	01A	OPAL $E_{cm}^{ee} = 88-94$ GeV
1738.1 ± 4.0	3.70M	ABREU	00F	DLPH $E_{cm}^{ee} = 88-94$ GeV
1751.1 ± 3.8	3.54M	ACCIARRI	00C	L3 $E_{cm}^{ee} = 88-94$ GeV
1744.0 ± 3.4	4.07M	BARATE	00C	ALEP $E_{cm}^{ee} = 88-94$ GeV

NODE=S044W5

Z BRANCHING RATIOS

NODE=S044220

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note "The Z boson" and ref. LEP-SLC 06).

NODE=S044220

$\Gamma(\text{hadrons})/\Gamma(e^+e^-)$

Γ_7/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
20.804 ± 0.050 OUR FIT				
20.902 ± 0.084	137.0K	¹ ABBIENDI	01A	OPAL $E_{cm}^{ee} = 88-94$ GeV
20.88 ± 0.12	117.8k	ABREU	00F	DLPH $E_{cm}^{ee} = 88-94$ GeV
20.816 ± 0.089	124.4k	ACCIARRI	00C	L3 $E_{cm}^{ee} = 88-94$ GeV
20.677 ± 0.075		² BARATE	00C	ALEP $E_{cm}^{ee} = 88-94$ GeV

NODE=S044R11

NODE=S044R11

• • • We do not use the following data for averages, fits, limits, etc. • • •

27.0 $^{+11.7}_{-8.8}$	12	³ ABRAMS	89D	MRK2 $E_{cm}^{ee} = 89-93$ GeV
------------------------	----	---------------------	-----	--------------------------------

¹ ABBIENDI 01A error includes approximately 0.067 due to statistics, 0.040 due to event selection systematics, 0.027 due to the theoretical uncertainty in t -channel prediction, and 0.014 due to LEP energy uncertainty.

NODE=S044R11;LINKAGE=DB

² BARATE 00C error includes approximately 0.062 due to statistics, 0.033 due to experimental systematics, and 0.026 due to the theoretical uncertainty in t -channel prediction.

NODE=S044R11;LINKAGE=AC

³ ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

NODE=S044R11;LINKAGE=AB

$\Gamma(\text{hadrons})/\Gamma(\mu^+\mu^-)$

Γ_7/Γ_2

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note "The Z boson" and ref. LEP-SLC 06).

NODE=S044R9

NODE=S044R9

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
20.785 ± 0.033 OUR FIT				
20.811 ± 0.058	182.8K	¹ ABBIENDI	01A	OPAL $E_{cm}^{ee} = 88-94$ GeV
20.65 ± 0.08	157.6k	ABREU	00F	DLPH $E_{cm}^{ee} = 88-94$ GeV
20.861 ± 0.097	113.4k	ACCIARRI	00C	L3 $E_{cm}^{ee} = 88-94$ GeV
20.799 ± 0.056		² BARATE	00C	ALEP $E_{cm}^{ee} = 88-94$ GeV

NODE=S044R9

• • • We do not use the following data for averages, fits, limits, etc. • • •

18.9 $^{+7.1}_{-5.3}$	13	³ ABRAMS	89D	MRK2 $E_{cm}^{ee} = 89-93$ GeV
-----------------------	----	---------------------	-----	--------------------------------

¹ ABBIENDI 01A error includes approximately 0.050 due to statistics and 0.027 due to event selection systematics.

NODE=S044R9;LINKAGE=DB

² BARATE 00C error includes approximately 0.053 due to statistics and 0.021 due to experimental systematics.

NODE=S044R9;LINKAGE=AC

³ ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

NODE=S044R9;LINKAGE=AB

$\Gamma(\text{hadrons})/\Gamma(\tau^+\tau^-)$ Γ_7/Γ_3

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06).

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
20.764±0.045 OUR FIT				
20.832±0.091	151.5K	¹ ABBIENDI	01A OPAL	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$
20.84 ±0.13	104.0k	ABREU	00F DLPH	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$
20.792±0.133	103.0k	ACCIARRI	00C L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$
20.707±0.062		² BARATE	00C ALEP	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

15.2 $\begin{smallmatrix} +4.8 \\ -3.9 \end{smallmatrix}$	21	³ ABRAMS	89D MRK2	$E_{\text{cm}}^{\text{ee}} = 89\text{--}93 \text{ GeV}$
---	----	---------------------	----------	---

¹ ABBIENDI 01A error includes approximately 0.055 due to statistics and 0.071 due to event selection systematics.

² BARATE 00C error includes approximately 0.054 due to statistics and 0.033 due to experimental systematics.

³ ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

 $\Gamma(\text{hadrons})/\Gamma(\ell^+\ell^-)$ Γ_7/Γ_4

ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

Our fit result is obtained requiring lepton universality.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
20.767±0.025 OUR FIT				
20.823±0.044	471.3K	¹ ABBIENDI	01A OPAL	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$
20.730±0.060	379.4k	ABREU	00F DLPH	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$
20.810±0.060	340.8k	ACCIARRI	00C L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$
20.725±0.039	500k	² BARATE	00C ALEP	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

18.9 $\begin{smallmatrix} +3.6 \\ -3.2 \end{smallmatrix}$	46	ABRAMS	89B MRK2	$E_{\text{cm}}^{\text{ee}} = 89\text{--}93 \text{ GeV}$
---	----	--------	----------	---

¹ ABBIENDI 01A error includes approximately 0.034 due to statistics and 0.027 due to event selection systematics.

² BARATE 00C error includes approximately 0.033 due to statistics, 0.020 due to experimental systematics, and 0.005 due to the theoretical uncertainty in t -channel prediction.

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_7/Γ

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

VALUE (%)	DOCUMENT ID
69.911±0.056 OUR FIT	

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_1/Γ

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

VALUE (%)	DOCUMENT ID
(3363.2±4.2) × 10⁻³ OUR FIT	

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_2/Γ

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

VALUE (%)	DOCUMENT ID
(3366.2±6.6) × 10⁻³ OUR FIT	

 $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_2/Γ_1

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

VALUE	DOCUMENT ID
1.0009±0.0028 OUR FIT	

 $\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$ Γ_3/Γ

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

VALUE (%)	DOCUMENT ID
(3369.6±8.3) × 10⁻³ OUR FIT	

NODE=S044R12

NODE=S044R12

NODE=S044R12

NODE=S044R12;LINKAGE=DB

NODE=S044R12;LINKAGE=AC

NODE=S044R12;LINKAGE=AB

NODE=S044R20

NODE=S044R20

NODE=S044R20

NODE=S044R20;LINKAGE=DB

NODE=S044R20;LINKAGE=AC

NODE=S044R8

NODE=S044R8

NODE=S044R8

NODE=S044R1

NODE=S044R1

NODE=S044R1

NODE=S044R2

NODE=S044R2

NODE=S044R2

NODE=S044R3

NODE=S044R3

NODE=S044R3

NODE=S044R10

NODE=S044R10

NODE=S044R10

$$\Gamma(\tau^+\tau^-)/\Gamma(e^+e^-)$$

 Γ_3/Γ_1

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

VALUE

DOCUMENT ID

1.0019 ± 0.0032 OUR FIT

NODE=S044R19

NODE=S044R19

NODE=S044R19

$$\Gamma(\ell^+\ell^-)/\Gamma_{\text{total}}$$

 Γ_4/Γ

ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

Our fit result assumes lepton universality.

This parameter is not directly used in the overall fit but is derived using the fit results; see the note “The Z boson” and ref. LEP-SLC 06.

VALUE (%)

DOCUMENT ID

(3365.8 ± 2.3) × 10⁻³ OUR FIT

NODE=S044R28

NODE=S044R28

NODE=S044R28

$$\Gamma(\ell^+\ell^-\ell^+\ell^-)/\Gamma_{\text{total}}$$

 Γ_5/Γ

Here ℓ indicates either e or μ .

VALUE (units 10⁻⁶)

EVTS

DOCUMENT ID

TECN

COMMENT

4.2^{+0.9}_{-0.8} ± 0.2

28

CHATRCHYAN 12BN CMS

 $E_{\text{cm}}^{pp} = 7 \text{ TeV}$

NODE=S044R01

NODE=S044R01

NODE=S044R01

$$\Gamma(\text{invisible})/\Gamma_{\text{total}}$$

 Γ_6/Γ

See the data, the note, and the fit result for the partial width, Γ_6 , above.

VALUE (%)

DOCUMENT ID

20.000 ± 0.055 OUR FIT

NODE=S044R30

NODE=S044R30

NODE=S044R30

$$\Gamma((u\bar{u}+c\bar{c})/2)/\Gamma(\text{hadrons})$$

 Γ_8/Γ_7

This quantity is the branching ratio of $Z \rightarrow$ “up-type” quarks to $Z \rightarrow$ hadrons. Except ACKERSTAFF 97T the values of $Z \rightarrow$ “up-type” and $Z \rightarrow$ “down-type” branchings are extracted from measurements of $\Gamma(\text{hadrons})$, and $\Gamma(Z \rightarrow \gamma + \text{jets})$ where γ is a high-energy (>5 or 7 GeV) isolated photon. As the experiments use different procedures and slightly different values of M_Z , $\Gamma(\text{hadrons})$ and α_s in their extraction procedures, our average has to be taken with caution.

VALUE

DOCUMENT ID

TECN

COMMENT

0.166 ± 0.009 OUR AVERAGE

 0.172^{+0.011}_{-0.010}

1 ABBIENDI

04E

OPAL

 $E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$

0.160 ± 0.019 ± 0.019

2 ACKERSTAFF

97T

OPAL

 $E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
 0.137^{+0.038}_{-0.054}

3 ABREU

95X

DLPH

 $E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

0.137 ± 0.033

4 ADRIANI

93

L3

 $E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$

¹ ABBIENDI 04E select photons with energy $> 7 \text{ GeV}$ and use $\Gamma(\text{hadrons}) = 1744.4 \pm 2.0 \text{ MeV}$ and $\alpha_s = 0.1172 \pm 0.002$ to obtain $\Gamma_u = 300^{+19}_{-18} \text{ MeV}$.

² ACKERSTAFF 97T measure $\Gamma_{u\bar{u}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}}) = 0.258 \pm 0.031 \pm 0.032$. To obtain this branching ratio authors use $R_c + R_b = 0.380 \pm 0.010$. This measurement is fully negatively correlated with the measurement of $\Gamma_{d\bar{d},s\bar{s}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}})$ given in the next data block.

³ ABREU 95X use $M_Z = 91.187 \pm 0.009 \text{ GeV}$, $\Gamma(\text{hadrons}) = 1725 \pm 12 \text{ MeV}$ and $\alpha_s = 0.123 \pm 0.005$. To obtain this branching ratio we divide their value of $C_{2/3} = 0.91^{+0.25}_{-0.36}$ by their value of $(3C_{1/3} + 2C_{2/3}) = 6.66 \pm 0.05$.

⁴ ADRIANI 93 use $M_Z = 91.181 \pm 0.022 \text{ GeV}$, $\Gamma(\text{hadrons}) = 1742 \pm 19 \text{ MeV}$ and $\alpha_s = 0.125 \pm 0.009$. To obtain this branching ratio we divide their value of $C_{2/3} = 0.92 \pm 0.22$ by their value of $(3C_{1/3} + 2C_{2/3}) = 6.720 \pm 0.076$.

NODE=S044R21

NODE=S044R21;LINKAGE=AB

NODE=S044R21;LINKAGE=D

NODE=S044R21;LINKAGE=C

NODE=S044R21;LINKAGE=B

$$\Gamma((d\bar{d}+s\bar{s}+b\bar{b})/3)/\Gamma(\text{hadrons})$$

 Γ_9/Γ_7

This quantity is the branching ratio of $Z \rightarrow$ “down-type” quarks to $Z \rightarrow$ hadrons. Except ACKERSTAFF 97T the values of $Z \rightarrow$ “up-type” and $Z \rightarrow$ “down-type” branchings are extracted from measurements of $\Gamma(\text{hadrons})$, and $\Gamma(Z \rightarrow \gamma + \text{jets})$ where γ is a high-energy (>5 or 7 GeV) isolated photon. As the experiments use different procedures and slightly different values of M_Z , $\Gamma(\text{hadrons})$ and α_s in their extraction procedures, our average has to be taken with caution.

VALUE

DOCUMENT ID

TECN

COMMENT

0.223 ± 0.006 OUR AVERAGE

0.218 ± 0.007

1 ABBIENDI

04E

OPAL

 $E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$

0.230 ± 0.010 ± 0.010

2 ACKERSTAFF

97T

OPAL

 $E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
 0.243^{+0.036}_{-0.026}

3 ABREU

95X

DLPH

 $E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

0.243 ± 0.022

4 ADRIANI

93

L3

 $E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$

NODE=S044R22

NODE=S044R22

NODE=S044R22

- ¹ ABBIENDI 04E select photons with energy > 7 GeV and use $\Gamma(\text{hadrons}) = 1744.4 \pm 2.0$ MeV and $\alpha_s = 0.1172 \pm 0.002$ to obtain $\Gamma_d = 381 \pm 12$ MeV.
- ² ACKERSTAFF 97T measure $\Gamma_{d\bar{d},s\bar{s}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}}) = 0.371 \pm 0.016 \pm 0.016$. To obtain this branching ratio authors use $R_c + R_b = 0.380 \pm 0.010$. This measurement is fully negatively correlated with the measurement of $\Gamma_{u\bar{u}}/(\Gamma_{d\bar{d}} + \Gamma_{u\bar{u}} + \Gamma_{s\bar{s}})$ presented in the previous data block.
- ³ ABREU 95X use $M_Z = 91.187 \pm 0.009$ GeV, $\Gamma(\text{hadrons}) = 1725 \pm 12$ MeV and $\alpha_s = 0.123 \pm 0.005$. To obtain this branching ratio we divide their value of $C_{1/3} = 1.62^{+0.24}_{-0.17}$ by their value of $(3C_{1/3} + 2C_{2/3}) = 6.66 \pm 0.05$.
- ⁴ ADRIANI 93 use $M_Z = 91.181 \pm 0.022$ GeV, $\Gamma(\text{hadrons}) = 1742 \pm 19$ MeV and $\alpha_s = 0.125 \pm 0.009$. To obtain this branching ratio we divide their value of $C_{1/3} = 1.63 \pm 0.15$ by their value of $(3C_{1/3} + 2C_{2/3}) = 6.720 \pm 0.076$.

NODE=S044R22;LINKAGE=AB

NODE=S044R22;LINKAGE=D

NODE=S044R22;LINKAGE=C

NODE=S044R22;LINKAGE=B

$R_c = \Gamma(c\bar{c})/\Gamma(\text{hadrons})$

 Γ_{10}/Γ_7

OUR FIT is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the note "The Z boson" and ref. LEP-SLC 06.

NODE=S044R29

NODE=S044R29

The Standard Model predicts $R_c = 0.1723$ for $m_t = 174.3$ GeV and $M_H = 150$ GeV.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1721\pm0.0030 OUR FIT			
0.1744 \pm 0.0031 \pm 0.0021	¹ ABE	05F SLD	$E_{cm}^{ee}=91.28$ GeV
0.1665 \pm 0.0051 \pm 0.0081	² ABREU	00 DLPH	$E_{cm}^{ee}=88-94$ GeV
0.1698 \pm 0.0069	³ BARATE	00B ALEP	$E_{cm}^{ee}=88-94$ GeV
0.180 \pm 0.011 \pm 0.013	⁴ ACKERSTAFF	98E OPAL	$E_{cm}^{ee}=88-94$ GeV
0.167 \pm 0.011 \pm 0.012	⁵ ALEXANDER	96R OPAL	$E_{cm}^{ee}=88-94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.1623 \pm 0.0085 \pm 0.0209	⁶ ABREU	95D DLPH	$E_{cm}^{ee}=88-94$ GeV

NODE=S044R29

¹ ABE 05F use hadronic Z decays collected during 1996–98 to obtain an enriched sample of $c\bar{c}$ events using a double tag method. The single c -tag is obtained with a neural network trained to perform flavor discrimination using as input several signatures (corrected secondary vertex mass, vertex decay length, multiplicity and total momentum of the hemisphere). A multitag approach is used, defining 4 regions of the output value of the neural network and R_c is extracted from a simultaneous fit to the count rates of the 4 different tags. The quoted systematic error includes an uncertainty of ± 0.0006 due to the uncertainty on R_b .

NODE=S044R29;LINKAGE=AB

² ABREU 00 obtain this result properly combining the measurement from the D^{*+} production rate ($R_c = 0.1610 \pm 0.0104 \pm 0.0077 \pm 0.0043$ (BR)) with that from the overall charm counting ($R_c = 0.1692 \pm 0.0047 \pm 0.0063 \pm 0.0074$ (BR)) in $c\bar{c}$ events. The systematic error includes an uncertainty of ± 0.0054 due to the uncertainty on the charmed hadron branching fractions.

NODE=S044R29;LINKAGE=Z

³ BARATE 00B use exclusive decay modes to independently determine the quantities $R_c \times f(c \rightarrow X)$, $X=D^0, D^+, D_s^+$, and Λ_c^+ . Estimating $R_c \times f(c \rightarrow \Xi_c/\Omega_c) = 0.0034$, they simply sum over all the charm decays to obtain $R_c = 0.1738 \pm 0.0047 \pm 0.0088 \pm 0.0075$ (BR). This is combined with all previous ALEPH measurements (BARATE 98T and BUSKULIC 94G, $R_c = 0.1681 \pm 0.0054 \pm 0.0062$) to obtain the quoted value.

NODE=S044R29;LINKAGE=T

⁴ ACKERSTAFF 98E use an inclusive/exclusive double tag. In one jet $D^{*\pm}$ mesons are exclusively reconstructed in several decay channels and in the opposite jet a slow pion (opposite charge inclusive $D^{*\pm}$) tag is used. The b content of this sample is measured by the simultaneous detection of a lepton in one jet and an inclusively reconstructed $D^{*\pm}$ meson in the opposite jet. The systematic error includes an uncertainty of ± 0.006 due to the external branching ratios.

NODE=S044R29;LINKAGE=H

⁵ ALEXANDER 96R obtain this value via direct charm counting, summing the partial contributions from D^0, D^+, D_s^+ , and Λ_c^+ , and assuming that strange-charmed baryons account for the 15% of the Λ_c^+ production. An uncertainty of ± 0.005 due to the uncertainties in the charm hadron branching ratios is included in the overall systematics.

NODE=S044R29;LINKAGE=G

⁶ ABREU 95D perform a maximum likelihood fit to the combined p and p_T distributions of single and dilepton samples. The second error includes an uncertainty of ± 0.0124 due to models and branching ratios.

NODE=S044R29;LINKAGE=AR

$R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$

 Γ_{11}/Γ_7

OUR FIT is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the note "The Z boson" and ref. LEP-SLC 06.

NODE=S044R18

NODE=S044R18

The Standard Model predicts $R_b = 0.21581$ for $m_t = 174.3$ GeV and $M_H = 150$ GeV.

VALUE	DOCUMENT ID	TECN	COMMENT
0.21629±0.00066 OUR FIT			
0.21594±0.00094±0.00075	¹ ABE	05F SLD	$E_{cm}^{ee}=91.28$ GeV
0.2174 ±0.0015 ±0.0028	² ACCIARRI	00 L3	$E_{cm}^{ee}=89-93$ GeV
0.2178 ±0.0011 ±0.0013	³ ABBIENDI	99B OPAL	$E_{cm}^{ee}=88-94$ GeV
0.21634±0.00067±0.00060	⁴ ABREU	99B DLPH	$E_{cm}^{ee}=88-94$ GeV
0.2159 ±0.0009 ±0.0011	⁵ BARATE	97F ALEP	$E_{cm}^{ee}=88-94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.2145 ±0.0089 ±0.0067	⁶ ABREU	95D DLPH	$E_{cm}^{ee}=88-94$ GeV
0.219 ±0.006 ±0.005	⁷ BUSKULIC	94G ALEP	$E_{cm}^{ee}=88-94$ GeV
0.251 ±0.049 ±0.030	⁸ JACOBSEN	91 MRK2	$E_{cm}^{ee}=91$ GeV

NODE=S044R18

OCCUR=2

NODE=S044R18;LINKAGE=AB

¹ ABE 05F use hadronic Z decays collected during 1996–98 to obtain an enriched sample of $b\bar{b}$ events using a double tag method. The single b -tag is obtained with a neural network trained to perform flavor discrimination using as input several signatures (corrected secondary vertex mass, vertex decay length, multiplicity and total momentum of the hemisphere; the key tag is obtained requiring the secondary vertex corrected mass to be above the D -meson mass). ABE 05F obtain $R_b = 0.21604 \pm 0.00098 \pm 0.00074$ where the systematic error includes an uncertainty of ± 0.00012 due to the uncertainty on R_c . The value reported here is obtained properly combining with ABE 98D. The quoted systematic error includes an uncertainty of ± 0.00012 due to the uncertainty on R_c .

² ACCIARRI 00 obtain this result using a double-tagging technique, with a high p_T lepton tag and an impact parameter tag in opposite hemispheres.

³ ABBIENDI 99B tag $Z \rightarrow b\bar{b}$ decays using leptons and/or separated decay vertices. The b -tagging efficiency is measured directly from the data using a double-tagging technique.

⁴ ABREU 99B obtain this result combining in a multivariate analysis several tagging methods (impact parameter and secondary vertex reconstruction, complemented by event shape variables). For R_c different from its Standard Model value of 0.172, R_b varies as $-0.024 \times (R_c - 0.172)$.

⁵ BARATE 97F combine the lifetime-mass hemisphere tag (BARATE 97E) with event shape information and lepton tag to identify $Z \rightarrow b\bar{b}$ candidates. They further use c - and uds -selection tags to identify the background. For R_c different from its Standard Model value of 0.172, R_b varies as $-0.019 \times (R_c - 0.172)$.

⁶ ABREU 95D perform a maximum likelihood fit to the combined p and p_T distributions of single and dilepton samples. The second error includes an uncertainty of ± 0.0023 due to models and branching ratios.

⁷ BUSKULIC 94G perform a simultaneous fit to the p and p_T spectra of both single and dilepton events.

⁸ JACOBSEN 91 tagged $b\bar{b}$ events by requiring coincidence of ≥ 3 tracks with significant impact parameters using vertex detector. Systematic error includes lifetime and decay uncertainties (± 0.014).

NODE=S044R18;LINKAGE=Z

NODE=S044R18;LINKAGE=IB

NODE=S044R18;LINKAGE=I

NODE=S044R18;LINKAGE=W

NODE=S044R18;LINKAGE=AR

NODE=S044R18;LINKAGE=BB

NODE=S044R18;LINKAGE=B

 $\Gamma(b\bar{b}b\bar{b})/\Gamma(\text{hadrons})$ **Γ_{12}/Γ_7**

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
5.2±1.9 OUR AVERAGE			
3.6±1.7±2.7	¹ ABBIENDI	01G OPAL	$E_{cm}^{ee}=88-94$ GeV
6.0±1.9±1.4	² ABREU	99U DLPH	$E_{cm}^{ee}=88-94$ GeV

NODE=S044B4
NODE=S044B4

¹ ABBIENDI 01G use a sample of four-jet events from hadronic Z decays. To enhance the $b\bar{b}b\bar{b}$ signal, at least three of the four jets are required to have a significantly detached secondary vertex.

² ABREU 99U force hadronic Z decays into 3 jets to use all the available phase space and require a b tag for every jet. This decay mode includes primary and secondary $4b$ production, e.g. from gluon splitting to $b\bar{b}$.

NODE=S044B4;LINKAGE=B

NODE=S044B4;LINKAGE=A

 $\Gamma(gg g)/\Gamma(\text{hadrons})$ **Γ_{13}/Γ_7**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6 × 10⁻²	95	¹ ABREU	96S DLPH	$E_{cm}^{ee}=88-94$ GeV

NODE=S044R63
NODE=S044R63

¹ This branching ratio is slightly dependent on the jet-finder algorithm. The value we quote is obtained using the JADE algorithm, while using the DURHAM algorithm ABREU 96S obtain an upper limit of 1.5×10^{-2} .

NODE=S044R63;LINKAGE=A

 $\Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}$ **Γ_{14}/Γ**

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<5.2 × 10⁻⁵	95	¹ ACCIARRI	95G L3	$E_{cm}^{ee}=88-94$ GeV
<5.5 × 10 ⁻⁵	95	ABREU	94B DLPH	$E_{cm}^{ee}=88-94$ GeV
<2.1 × 10 ⁻⁴	95	DECAMP	92 ALEP	$E_{cm}^{ee}=88-94$ GeV
<1.4 × 10 ⁻⁴	95	AKRAWY	91F OPAL	$E_{cm}^{ee}=88-94$ GeV

NODE=S044R13
NODE=S044R13

¹ This limit is for both decay modes $Z \rightarrow \pi^0 \gamma / \gamma \gamma$ which are indistinguishable in ACCIARRI 95G.

NODE=S044R13;LINKAGE=C

$\Gamma(\eta\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.6 \times 10^{-5}$	95	ACCIARRI	95G L3	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<8.0 \times 10^{-5}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<5.1 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<2.0 \times 10^{-4}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{15}/Γ

NODE=S044R14
NODE=S044R14

 $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.5 \times 10^{-4}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{16}/Γ

NODE=S044R48
NODE=S044R48

 $\Gamma(\eta'(958)\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.2 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{17}/Γ

NODE=S044R15
NODE=S044R15

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

This decay would violate the Landau-Yang theorem.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-5}$	95	¹ ACCIARRI	95G L3	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<5.5 \times 10^{-5}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<1.4 \times 10^{-4}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{18}/Γ

NODE=S044R16
NODE=S044R16
NODE=S044R16

¹ This limit is for both decay modes $Z \rightarrow \pi^0 \gamma / \gamma \gamma$ which are indistinguishable in ACCIA-RRI 95G.

NODE=S044R16;LINKAGE=C

 $\Gamma(\gamma\gamma\gamma)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.0 \times 10^{-5}$	95	¹ ACCIARRI	95C L3	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<1.7 \times 10^{-5}$	95	¹ ABREU	94B DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<6.6 \times 10^{-5}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{19}/Γ

NODE=S044R17
NODE=S044R17

¹ Limit derived in the context of composite Z model.

NODE=S044R17;LINKAGE=AB

 $\Gamma(\pi^\pm W^\mp)/\Gamma_{\text{total}}$

The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{20}/Γ

NODE=S044R32
NODE=S044R32
NODE=S044R32

 $\Gamma(\rho^\pm W^\mp)/\Gamma_{\text{total}}$

The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.3 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{21}/Γ

NODE=S044R33
NODE=S044R33
NODE=S044R33

 $\Gamma(J/\psi(1S)X)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

 Γ_{22}/Γ

NODE=S044R23
NODE=S044R23

$3.51^{+0.23}_{-0.25}$ OUR AVERAGE Error includes scale factor of 1.1.

$3.21 \pm 0.21^{+0.19}_{-0.28}$	553	¹ ACCIARRI	99F L3	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$3.9 \pm 0.2 \pm 0.3$	511	² ALEXANDER	96B OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$3.73 \pm 0.39 \pm 0.36$	153	³ ABREU	94P DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

¹ ACCIARRI 99F combine $\mu^+ \mu^-$ and $e^+ e^- J/\psi(1S)$ decay channels. The branching ratio for prompt $J/\psi(1S)$ production is measured to be $(2.1 \pm 0.6 \pm 0.4^{+0.4}_{-0.2}(\text{theor.})) \times 10^{-4}$.

NODE=S044R23;LINKAGE=F

² ALEXANDER 96B identify $J/\psi(1S)$ from the decays into lepton pairs. $(4.8 \pm 2.4)\%$ of this branching ratio is due to prompt $J/\psi(1S)$ production (ALEXANDER 96N).

NODE=S044R23;LINKAGE=D

³ Combining $\mu^+ \mu^-$ and $e^+ e^-$ channels and taking into account the common systematic errors. $(7.7^{+6.3}_{-5.4})\%$ of this branching ratio is due to prompt $J/\psi(1S)$ production.

NODE=S044R23;LINKAGE=C

 $\Gamma(\psi(2S)X)/\Gamma_{\text{total}}$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.60 ± 0.29 OUR AVERAGE				
$1.6 \pm 0.5 \pm 0.3$	39	¹ ACCIARRI	97J L3	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$1.6 \pm 0.3 \pm 0.2$	46.9	² ALEXANDER	96B OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$1.60 \pm 0.73 \pm 0.33$	5.4	³ ABREU	94P DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{23}/Γ

NODE=S044R60
NODE=S044R60

- ¹ ACCIARRI 97J measure this branching ratio via the decay channel $\psi(2S) \rightarrow \ell^+ \ell^-$ ($\ell = \mu, e$).
- ² ALEXANDER 96B measure this branching ratio via the decay channel $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$, with $J/\psi \rightarrow \ell^+ \ell^-$.
- ³ ABREU 94P measure this branching ratio via decay channel $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$, with $J/\psi \rightarrow \mu^+ \mu^-$.

NODE=S044R60;LINKAGE=C

NODE=S044R60;LINKAGE=BB

NODE=S044R60;LINKAGE=A

 $\Gamma(\chi_{c1}(1P)X)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.9±0.7 OUR AVERAGE				
2.7±0.6±0.5	33	¹ ACCIARRI	97J L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV
5.0±2.1 ^{+1.5} _{-0.9}	6.4	² ABREU	94P DLPH	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

NODE=S044R42
NODE=S044R42

- ¹ ACCIARRI 97J measure this branching ratio via the decay channel $\chi_{c1} \rightarrow J/\psi + \gamma$, with $J/\psi \rightarrow \ell^+ \ell^-$ ($\ell = \mu, e$). The $M(\ell^+ \ell^- \gamma) - M(\ell^+ \ell^-)$ mass difference spectrum is fitted with two gaussian shapes for χ_{c1} and χ_{c2} .

NODE=S044R42;LINKAGE=C

- ² This branching ratio is measured via the decay channel $\chi_{c1} \rightarrow J/\psi + \gamma$, with $J/\psi \rightarrow \mu^+ \mu^-$.

NODE=S044R42;LINKAGE=A

 $\Gamma(\chi_{c2}(1P)X)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.2 × 10⁻³	90	¹ ACCIARRI	97J L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

NODE=S044R65
NODE=S044R65

- ¹ ACCIARRI 97J derive this limit via the decay channel $\chi_{c2} \rightarrow J/\psi + \gamma$, with $J/\psi \rightarrow \ell^+ \ell^-$ ($\ell = \mu, e$). The $M(\ell^+ \ell^- \gamma) - M(\ell^+ \ell^-)$ mass difference spectrum is fitted with two gaussian shapes for χ_{c1} and χ_{c2} .

NODE=S044R65;LINKAGE=A

 $\Gamma(\Upsilon(1S)X + \Upsilon(2S)X + \Upsilon(3S)X)/\Gamma_{\text{total}}$ $\Gamma_{26}/\Gamma = (\Gamma_{27} + \Gamma_{28} + \Gamma_{29})/\Gamma$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.0±0.4±0.22	6.4	¹ ALEXANDER	96F OPAL	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

NODE=S044R62
NODE=S044R62

- ¹ ALEXANDER 96F identify the Υ (which refers to any of the three lowest bound states) through its decay into $e^+ e^-$ and $\mu^+ \mu^-$. The systematic error includes an uncertainty of ± 0.2 due to the production mechanism.

NODE=S044R62;LINKAGE=A

 $\Gamma(\Upsilon(1S)X)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.4 × 10⁻⁵	95	¹ ACCIARRI	99F L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

NODE=S044R66
NODE=S044R66

- ¹ ACCIARRI 99F search for $\Upsilon(1S)$ through its decay into $\ell^+ \ell^-$ ($\ell = e$ or μ).

NODE=S044R66;LINKAGE=B

 $\Gamma(\Upsilon(2S)X)/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<13.9 × 10⁻⁵	95	¹ ACCIARRI	97R L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

NODE=S044R67
NODE=S044R67

- ¹ ACCIARRI 97R search for $\Upsilon(2S)$ through its decay into $\ell^+ \ell^-$ ($\ell = e$ or μ).

NODE=S044R67;LINKAGE=A

 $\Gamma(\Upsilon(3S)X)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9.4 × 10⁻⁵	95	¹ ACCIARRI	97R L3	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

NODE=S044R68
NODE=S044R68

- ¹ ACCIARRI 97R search for $\Upsilon(3S)$ through its decay into $\ell^+ \ell^-$ ($\ell = e$ or μ).

NODE=S044R68;LINKAGE=A

 $\Gamma((D^0/\bar{D}^0)X)/\Gamma(\text{hadrons})$ Γ_{30}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.296±0.019±0.021	369	¹ ABREU	93I DLPH	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

NODE=S044R43
NODE=S044R43

- ¹ The (D^0/\bar{D}^0) states in ABREU 93I are detected by the $K\pi$ decay mode. This is a corrected result (see the erratum of ABREU 93I).

NODE=S044R43;LINKAGE=A

 $\Gamma(D^\pm X)/\Gamma(\text{hadrons})$ Γ_{31}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.174±0.016±0.018	539	¹ ABREU	93I DLPH	$E_{\text{cm}}^{\text{ee}} = 88\text{--}94$ GeV

NODE=S044R44
NODE=S044R44

- ¹ The D^\pm states in ABREU 93I are detected by the $K\pi\pi$ decay mode. This is a corrected result (see the erratum of ABREU 93I).

NODE=S044R44;LINKAGE=A

$\Gamma(D^*(2010)^\pm X)/\Gamma(\text{hadrons})$ Γ_{32}/Γ_7

The value is for the sum of the charge states indicated.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.163±0.019 OUR AVERAGE		Error includes scale factor of 1.3.		
0.155±0.010±0.013	358	¹ ABREU	93I DLPH	$E_{cm}^{ee} = 88-94$ GeV
0.21 ±0.04	362	² DECAMP	91J ALEP	$E_{cm}^{ee} = 88-94$ GeV

¹ $D^*(2010)^\pm$ in ABREU 93I are reconstructed from $D^0\pi^\pm$, with $D^0 \rightarrow K^-\pi^+$. The new CLEO II measurement of $B(D^{*\pm} \rightarrow D^0\pi^\pm) = (68.1 \pm 1.6)\%$ is used. This is a corrected result (see the erratum of ABREU 93I).

² DECAMP 91J report $B(D^*(2010)^+ \rightarrow D^0\pi^+) B(D^0 \rightarrow K^-\pi^+) \Gamma(D^*(2010)^\pm X) / \Gamma(\text{hadrons}) = (5.11 \pm 0.34) \times 10^{-3}$. They obtained the above number assuming $B(D^0 \rightarrow K^-\pi^+) = (3.62 \pm 0.34 \pm 0.44)\%$ and $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (55 \pm 4)\%$. We have rescaled their original result of 0.26 ± 0.05 taking into account the new CLEO II branching ratio $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.6)\%$.

NODE=S044R24

NODE=S044R24

NODE=S044R24

NODE=S044R24;LINKAGE=B

NODE=S044R24;LINKAGE=A

 $\Gamma(D_{s1}(2536)^\pm X)/\Gamma(\text{hadrons})$ Γ_{33}/Γ_7 $D_{s1}(2536)^\pm$ is an expected orbitally-excited state of the D_s meson.

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
0.52±0.09±0.06	92	¹ HEISTER	02B ALEP	$E_{cm}^{ee} = 88-94$ GeV

¹ HEISTER 02B reconstruct this meson in the decay modes $D_{s1}(2536)^\pm \rightarrow D^{*\pm}K^0$ and $D_{s1}(2536)^\pm \rightarrow D^{*0}K^\pm$. The quoted branching ratio assumes that the decay width of the $D_{s1}(2536)$ is saturated by the two measured decay modes.

NODE=S044R70

NODE=S044R70

NODE=S044R70

NODE=S044R70;LINKAGE=A

 $\Gamma(D_{sJ}(2573)^\pm X)/\Gamma(\text{hadrons})$ Γ_{34}/Γ_7 $D_{sJ}(2573)^\pm$ is an expected orbitally-excited state of the D_s meson.

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
0.83±0.29^{+0.07}_{-0.13}	64	¹ HEISTER	02B ALEP	$E_{cm}^{ee} = 88-94$ GeV

¹ HEISTER 02B reconstruct this meson in the decay mode $D_{s2}^*(2573)^\pm \rightarrow D^0K^\pm$. The quoted branching ratio assumes that the detected decay mode represents 45% of the full decay width.

NODE=S044R71

NODE=S044R71

NODE=S044R71

NODE=S044R71;LINKAGE=A

 $\Gamma(D^{*'}(2629)^\pm X)/\Gamma(\text{hadrons})$ Γ_{35}/Γ_7 $D^{*'}(2629)^\pm$ is a predicted radial excitation of the $D^*(2010)^\pm$ meson.

VALUE	DOCUMENT ID	TECN	COMMENT
searched for	¹ ABBIENDI	01N OPAL	$E_{cm}^{ee} = 88-94$ GeV

¹ ABBIENDI 01N searched for the decay mode $D^{*'}(2629)^\pm \rightarrow D^{*\pm}\pi^+\pi^-$ with $D^{*+} \rightarrow D^0\pi^+$, and $D^0 \rightarrow K^-\pi^+$. They quote a 95% CL limit for $Z \rightarrow D^{*'}(2629)^\pm \times B(D^{*'}(2629)^+ \rightarrow D^{*+}\pi^+\pi^-) < 3.1 \times 10^{-3}$.

NODE=S044R69

NODE=S044R69

NODE=S044R69

NODE=S044R69;LINKAGE=A

 $\Gamma(B^*X)/[\Gamma(BX) + \Gamma(B^*X)]$ $\Gamma_{37}/(\Gamma_{36}+\Gamma_{37})$ As the experiments assume different values of the b -baryon contribution, our average should be taken with caution.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.75 ±0.04 OUR AVERAGE				
0.760±0.036±0.083		¹ ACKERSTAFF	97M OPAL	$E_{cm}^{ee} = 88-94$ GeV
0.771±0.026±0.070		² BUSKULIC	96D ALEP	$E_{cm}^{ee} = 88-94$ GeV
0.72 ±0.03 ±0.06		³ ABREU	95R DLPH	$E_{cm}^{ee} = 88-94$ GeV
0.76 ±0.08 ±0.06	1378	⁴ ACCIARRI	95B L3	$E_{cm}^{ee} = 88-94$ GeV

¹ ACKERSTAFF 97M use an inclusive B reconstruction method and assume a $(13.2 \pm 4.1)\%$ b -baryon contribution. The value refers to a b -flavored meson mixture of B_u , B_d , and B_s .

² BUSKULIC 96D use an inclusive reconstruction of B hadrons and assume a $(12.2 \pm 4.3)\%$ b -baryon contribution. The value refers to a b -flavored mixture of B_u , B_d , and B_s .

³ ABREU 95R use an inclusive B -reconstruction method and assume a $(10 \pm 4)\%$ b -baryon contribution. The value refers to a b -flavored meson mixture of B_u , B_d , and B_s .

⁴ ACCIARRI 95B assume a 9.4% b -baryon contribution. The value refers to a b -flavored mixture of B_u , B_d , and B_s .

NODE=S044R61

NODE=S044R61

NODE=S044R61

NODE=S044R61;LINKAGE=D

NODE=S044R61;LINKAGE=C

NODE=S044R61;LINKAGE=B

NODE=S044R61;LINKAGE=A

$\Gamma(B^+X)/\Gamma(\text{hadrons})$ Γ_{38}/Γ_7

"OUR EVALUATION" is obtained using our current values for $f(\bar{b} \rightarrow B^+)$ and $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$. We calculate $\Gamma(B^+X)/\Gamma(\text{hadrons}) = R_b \times f(\bar{b} \rightarrow B^+)$. The decay fraction $f(\bar{b} \rightarrow B^+)$ was provided by the Heavy Flavor Averaging Group (HFAG, <http://www.slac.stanford.edu/xorg/hfag/osc/PDG.2009/#FRACZ>).

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0869 ± 0.0019 OUR EVALUATION**0.0887 ± 0.0030**¹ ABDALLAH 03K DLPH $E_{cm}^{ee} = 88-94$ GeV

¹ ABDALLAH 03K measure the production fraction of B^+ mesons in hadronic Z decays $f(B^+) = (40.99 \pm 0.82 \pm 1.11)\%$. The value quoted here is obtained multiplying this production fraction by our value of $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$.

NODE=S044B+X

NODE=S044B+X

NODE=S044B+X

→ UNCHECKED ←

NODE=S044B+X;LINKAGE=AB

 $\Gamma(B_s^0X)/\Gamma(\text{hadrons})$ Γ_{39}/Γ_7

"OUR EVALUATION" is obtained using our current values for $f(\bar{b} \rightarrow B_s^0)$ and $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$. We calculate $\Gamma(B_s^0X)/\Gamma(\text{hadrons}) = R_b \times f(\bar{b} \rightarrow B_s^0)$. The decay fraction $f(\bar{b} \rightarrow B_s^0)$ was provided by the Heavy Flavor Averaging Group (HFAG, <http://www.slac.stanford.edu/xorg/hfag/osc/PDG.2009/#FRACZ>).

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

0.0227 ± 0.0019 OUR EVALUATION

seen

¹ ABREU 92M DLPH $E_{cm}^{ee} = 88-94$ GeV

seen

² ACTON 92N OPAL $E_{cm}^{ee} = 88-94$ GeV

seen

³ BUSKULIC 92E ALEP $E_{cm}^{ee} = 88-94$ GeV

¹ ABREU 92M reported value is $\Gamma(B_s^0X) \times B(B_s^0 \rightarrow D_s \mu \nu_\mu X) \times B(D_s \rightarrow \phi \pi)/\Gamma(\text{hadrons}) = (18 \pm 8) \times 10^{-5}$.

² ACTON 92N find evidence for B_s^0 production using D_s - ℓ correlations, with $D_s^+ \rightarrow \phi \pi^+$ and $K^*(892)K^+$. Assuming R_b from the Standard Model and averaging over the e and μ channels, authors measure the product branching fraction to be $f(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell X) \times B(D_s^- \rightarrow \phi \pi^-) = (3.9 \pm 1.1 \pm 0.8) \times 10^{-4}$.

³ BUSKULIC 92E find evidence for B_s^0 production using D_s - ℓ correlations, with $D_s^+ \rightarrow \phi \pi^+$ and $K^*(892)K^+$. Using $B(D_s^+ \rightarrow \phi \pi^+) = (2.7 \pm 0.7)\%$ and summing up the e and μ channels, the weighted average product branching fraction is measured to be $B(\bar{b} \rightarrow B_s^0) \times B(B_s^0 \rightarrow D_s^- \ell^+ \nu_\ell X) = 0.040 \pm 0.011^{+0.010}_{-0.012}$.

NODE=S044R49

NODE=S044R49

NODE=S044R49

→ UNCHECKED ←

NODE=S044R49;LINKAGE=A

NODE=S044R49;LINKAGE=C

NODE=S044R49;LINKAGE=B

 $\Gamma(B_c^+X)/\Gamma(\text{hadrons})$ Γ_{40}/Γ_7

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

searched for

¹ ACKERSTAFF 98O OPAL $E_{cm}^{ee} = 88-94$ GeV

searched for

² ABREU 97E DLPH $E_{cm}^{ee} = 88-94$ GeV

searched for

³ BARATE 97H ALEP $E_{cm}^{ee} = 88-94$ GeV

¹ ACKERSTAFF 98O searched for the decay modes $B_c \rightarrow J/\psi \pi^+$, $J/\psi a_1^+$, and $J/\psi \ell^+ \nu_\ell$, with $J/\psi \rightarrow \ell^+ \ell^-$, $\ell = e, \mu$. The number of candidates (background) for the three decay modes is 2 (0.63 ± 0.2), 0 (1.10 ± 0.22), and 1 (0.82 ± 0.19) respectively. Interpreting the $2 B_c \rightarrow J/\psi \pi^+$ candidates as signal, they report $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi \pi^+)/\Gamma(\text{hadrons}) = (3.8^{+5.0}_{-2.4} \pm 0.5) \times 10^{-5}$. Interpreted as background, the 90% CL bounds are $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi \pi^+)/\Gamma(\text{hadrons}) < 1.06 \times 10^{-4}$, $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi a_1^+)/\Gamma(\text{hadrons}) < 5.29 \times 10^{-4}$, $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi \ell^+ \nu_\ell)/\Gamma(\text{hadrons}) < 6.96 \times 10^{-5}$.

² ABREU 97E searched for the decay modes $B_c \rightarrow J/\psi \pi^+$, $J/\psi \ell^+ \nu_\ell$, and $J/\psi (3\pi)^+$, with $J/\psi \rightarrow \ell^+ \ell^-$, $\ell = e, \mu$. The number of candidates (background) for the three decay modes is 1 (1.7), 0 (0.3), and 1 (2.3) respectively. They report the following 90% CL limits: $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi \pi^+)/\Gamma(\text{hadrons}) < (1.05-0.84) \times 10^{-4}$, $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi \ell \nu_\ell)/\Gamma(\text{hadrons}) < (5.8-5.0) \times 10^{-5}$, $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi (3\pi)^+)/\Gamma(\text{hadrons}) < 1.75 \times 10^{-4}$, where the ranges are due to the predicted B_c lifetime (0.4-1.4) ps.

³ BARATE 97H searched for the decay modes $B_c \rightarrow J/\psi \pi^+$ and $J/\psi \ell^+ \nu_\ell$ with $J/\psi \rightarrow \ell^+ \ell^-$, $\ell = e, \mu$. The number of candidates (background) for the two decay modes is 0 (0.44) and 2 (0.81) respectively. They report the following 90% CL limits: $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi \pi^+)/\Gamma(\text{hadrons}) < 3.6 \times 10^{-5}$ and $\Gamma(B_c^+X) \times B(B_c \rightarrow J/\psi \ell^+ \nu_\ell)/\Gamma(\text{hadrons}) < 5.2 \times 10^{-5}$.

NODE=S044BCC
NODE=S044BCC

NODE=S044BCC;LINKAGE=A

NODE=S044BCC;LINKAGE=B

NODE=S044BCC;LINKAGE=C

$\Gamma(\Lambda_c^+ X)/\Gamma(\text{hadrons})$ Γ_{41}/Γ_7

VALUE

DOCUMENT ID

TECN

COMMENT

0.022 ± 0.005 OUR AVERAGE

0.024 ± 0.005 ± 0.006

¹ ALEXANDER 96R OPAL $E_{cm}^{ee} = 88-94$ GeV

0.021 ± 0.003 ± 0.005

² BUSKULIC 96Y ALEP $E_{cm}^{ee} = 88-94$ GeV

¹ ALEXANDER 96R measure $R_b \times f(b \rightarrow \Lambda_c^+ X) \times B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (0.122 \pm 0.023 \pm 0.010)\%$ in hadronic Z decays; the value quoted here is obtained using our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3)\%$. The first error is the total experiment's error and the second error is the systematic error due to the branching fraction uncertainty.

² BUSKULIC 96Y obtain the production fraction of Λ_c^+ baryons in hadronic Z decays $f(b \rightarrow \Lambda_c^+ X) = 0.110 \pm 0.014 \pm 0.006$ using $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (4.4 \pm 0.6)\%$; we have rescaled using our best value $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (5.0 \pm 1.3)\%$ obtaining $f(b \rightarrow \Lambda_c^+ X) = 0.097 \pm 0.013 \pm 0.025$ where the first error is their total experiment's error and the second error is the systematic error due to the branching fraction uncertainty. The value quoted here is obtained multiplying this production fraction by our value of $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$.

NODE=S044LCX
NODE=S044LCX

NODE=S044LCX;LINKAGE=AL

NODE=S044LCX;LINKAGE=BU

 $\Gamma(\Xi_c^0 X)/\Gamma(\text{hadrons})$ Γ_{42}/Γ_7

VALUE

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen

¹ ABDALLAH 05C DLPH $E_{cm}^{ee} = 88-94$ GeV

¹ ABDALLAH 05C searched for the charmed strange baryon Ξ_c^0 in the decay channel $\Xi_c^0 \rightarrow \Xi^- \pi^+ (\Xi^- \rightarrow \Lambda \pi^-)$. The production rate is measured to be $f_{\Xi_c^0} \times B(\Xi_c^0 \rightarrow \Xi^- \pi^+) = (4.7 \pm 1.4 \pm 1.1) \times 10^{-4}$ per hadronic Z decay.

NODE=S044XIC
NODE=S044XIC

NODE=S044XIC;LINKAGE=AB

 $\Gamma(\Xi_b X)/\Gamma(\text{hadrons})$ Γ_{43}/Γ_7 Here Ξ_b is used as a notation for the strange b -baryon states Ξ_b^- and Ξ_b^0 .

VALUE

DOCUMENT ID

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen

¹ ABDALLAH 05C DLPH $E_{cm}^{ee} = 88-94$ GeV

seen

² BUSKULIC 96T ALEP $E_{cm}^{ee} = 88-94$ GeV

seen

³ ABREU 95V DLPH $E_{cm}^{ee} = 88-94$ GeV

¹ ABDALLAH 05C searched for the beauty strange baryon Ξ_b in the inclusive semileptonic decay channel $\Xi_b \rightarrow \Xi^- \ell^- \bar{\nu}_\ell X$. Evidence for the Ξ_b production is seen from the observation of Ξ^\mp production accompanied by a lepton of the same sign. From the excess of "right-sign" pairs $\Xi^\mp \ell^\mp$ compared to "wrong-sign" pairs $\Xi^\mp \ell^\pm$ the production rate is measured to be $B(b \rightarrow \Xi_b) \times B(\Xi_b \rightarrow \Xi^- \ell^- X) = (3.0 \pm 1.0 \pm 0.3) \times 10^{-4}$ per lepton species, averaged over electrons and muons.

² BUSKULIC 96T investigate Ξ -lepton correlations and find a significant excess of "right-sign" pairs $\Xi^\mp \ell^\mp$ compared to "wrong-sign" pairs $\Xi^\mp \ell^\pm$. This excess is interpreted as evidence for Ξ_b semileptonic decay. The measured product branching ratio is $B(b \rightarrow \Xi_b) \times B(\Xi_b \rightarrow X_c X \ell^- \bar{\nu}_\ell) \times B(X_c \rightarrow \Xi^- X') = (5.4 \pm 1.1 \pm 0.8) \times 10^{-4}$ per lepton species, averaged over electrons and muons, with X_c a charmed baryon.

³ ABREU 95V observe an excess of "right-sign" pairs $\Xi^\mp \ell^\mp$ compared to "wrong-sign" pairs $\Xi^\mp \ell^\pm$ in jets: this excess is interpreted as evidence for the beauty strange baryon Ξ_b production, with $\Xi_b \rightarrow \Xi^- \ell^- \bar{\nu}_\ell X$. They find that the probability for this signal to come from non b -baryon decays is less than 5×10^{-4} and that Λ_b decays can account for less than 10% of these events. The Ξ_b production rate is then measured to be $B(b \rightarrow \Xi_b) \times B(\Xi_b \rightarrow \Xi^- \ell^- X) = (5.9 \pm 2.1 \pm 1.0) \times 10^{-4}$ per lepton species, averaged over electrons and muons.

NODE=S044XIB

NODE=S044XIB
NODE=S044XIB

NODE=S044XIB;LINKAGE=AB

NODE=S044XIB;LINKAGE=BU

NODE=S044XIB;LINKAGE=AR

 $\Gamma(b\text{-baryon } X)/\Gamma(\text{hadrons})$ Γ_{44}/Γ_7

"OUR EVALUATION" is obtained using our current values for $f(b \rightarrow b\text{-baryon})$ and $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$. We calculate $\Gamma(b\text{-baryon } X)/\Gamma(\text{hadrons}) = R_b \times f(b \rightarrow b\text{-baryon})$. The decay fraction $f(b \rightarrow b\text{-baryon})$ was provided by the Heavy Flavor Averaging Group (HFAG, <http://www.slac.stanford.edu/xorg/hfag/osc/PDG.2009>).

VALUE

DOCUMENT ID

TECN

COMMENT

0.0197 ± 0.0032 OUR EVALUATION**0.0221 ± 0.0015 ± 0.0058**¹ BARATE 98V ALEP $E_{cm}^{ee} = 88-94$ GeV

¹ BARATE 98V use the overall number of identified protons in b -hadron decays to measure $f(b \rightarrow b\text{-baryon}) = 0.102 \pm 0.007 \pm 0.027$. They assume $BR(b\text{-baryon} \rightarrow p X) = (58 \pm 6)\%$ and $BR(B_s^0 \rightarrow p X) = (8.0 \pm 4.0)\%$. The value quoted here is obtained multiplying this production fraction by our value of $R_b = \Gamma(b\bar{b})/\Gamma(\text{hadrons})$.

NODE=S044BBR

NODE=S044BBR

NODE=S044BBR
→ UNCHECKED ←

NODE=S044BBR;LINKAGE=BA

$\Gamma(\text{anomalous } \gamma + \text{hadrons})/\Gamma_{\text{total}}$ Γ_{45}/Γ

Limits on additional sources of prompt photons beyond expectations for final-state bremsstrahlung.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-3}$	95	¹ AKRAWY 90J	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

¹ AKRAWY 90J report $\Gamma(\gamma X) < 8.2 \text{ MeV}$ at 95%CL. They assume a three-body $\gamma q \bar{q}$ distribution and use $E(\gamma) > 10 \text{ GeV}$.

NODE=S044R31

NODE=S044R31

NODE=S044R31

NODE=S044R31;LINKAGE=A

 $\Gamma(e^+ e^- \gamma)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-4}$	95	¹ ACTON 91B	OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$

¹ ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy ($> 0.9 \text{ GeV}$).

NODE=S044R34

NODE=S044R34

NODE=S044R34;LINKAGE=A

 $\Gamma(\mu^+ \mu^- \gamma)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-4}$	95	¹ ACTON 91B	OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$

¹ ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy ($> 0.9 \text{ GeV}$).

NODE=S044R35

NODE=S044R35

NODE=S044R35;LINKAGE=A

 $\Gamma(\tau^+ \tau^- \gamma)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.3 \times 10^{-4}$	95	¹ ACTON 91B	OPAL	$E_{\text{cm}}^{ee} = 91.2 \text{ GeV}$

¹ ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy ($> 0.9 \text{ GeV}$).

NODE=S044R36

NODE=S044R36

NODE=S044R36;LINKAGE=A

 $\Gamma(\ell^+ \ell^- \gamma \gamma)/\Gamma_{\text{total}}$ Γ_{49}/Γ

The value is the sum over $\ell = e, \mu, \tau$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.8 \times 10^{-6}$	95	¹ ACTON 93E	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

¹ For $m_{\gamma\gamma} = 60 \pm 5 \text{ GeV}$.

NODE=S044R45

NODE=S044R45

NODE=S044R45

NODE=S044R45;LINKAGE=A

 $\Gamma(q \bar{q} \gamma \gamma)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.5 \times 10^{-6}$	95	¹ ACTON 93E	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

¹ For $m_{\gamma\gamma} = 60 \pm 5 \text{ GeV}$.

NODE=S044R46

NODE=S044R46

NODE=S044R46;LINKAGE=A

 $\Gamma(\nu \bar{\nu} \gamma \gamma)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.1 \times 10^{-6}$	95	¹ ACTON 93E	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

¹ For $m_{\gamma\gamma} = 60 \pm 5 \text{ GeV}$.

NODE=S044R47

NODE=S044R47

NODE=S044R47;LINKAGE=A

 $\Gamma(e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{52}/Γ

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-6}$	95	ABREU 97C	DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
$<1.7 \times 10^{-6}$	95	AKERS 95W	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
$<0.6 \times 10^{-5}$	95	ADRIANI 93I	L3	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
$<2.6 \times 10^{-5}$	95	DECAMP 92	ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

NODE=S044R27

NODE=S044R27

NODE=S044R27

 $\Gamma(e^\pm \mu^\mp)/\Gamma(e^+ e^-)$ Γ_{52}/Γ_1

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.07	90	ALBAJAR 89	UA1	$E_{\text{cm}}^{p\bar{p}} = 546,630 \text{ GeV}$

NODE=S044R6

NODE=S044R6

NODE=S044R6

 $\Gamma(e^\pm \tau^\mp)/\Gamma_{\text{total}}$ Γ_{53}/Γ

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.2 \times 10^{-5}$	95	ABREU 97C	DLPH	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
$<9.8 \times 10^{-6}$	95	AKERS 95W	OPAL	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
$<1.3 \times 10^{-5}$	95	ADRIANI 93I	L3	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$
$<1.2 \times 10^{-4}$	95	DECAMP 92	ALEP	$E_{\text{cm}}^{ee} = 88\text{--}94 \text{ GeV}$

NODE=S044R25

NODE=S044R25

NODE=S044R25

$\Gamma(\mu^\pm \tau^\mp)/\Gamma_{\text{total}}$

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-5}$	95	ABREU	97C DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<1.7 \times 10^{-5}$	95	AKERS	95W OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<1.9 \times 10^{-5}$	95	ADRIANI	93I L3	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$
$<1.0 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

 Γ_{54}/Γ

NODE=S044R26

NODE=S044R26

NODE=S044R26

 $\Gamma(pe)/\Gamma_{\text{total}}$

Test of baryon number and lepton number conservations. Charge conjugate states are implied.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	95	¹ ABBIENDI	99I OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

¹ ABBIENDI 99I give the 95%CL limit on the partial width $\Gamma(Z^0 \rightarrow pe) < 4.6 \text{ KeV}$ and we have transformed it into a branching ratio.

 Γ_{55}/Γ

NODE=S044PE

NODE=S044PE

NODE=S044PE

NODE=S044PE;LINKAGE=A

 $\Gamma(p\mu)/\Gamma_{\text{total}}$

Test of baryon number and lepton number conservations. Charge conjugate states are implied.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-6}$	95	¹ ABBIENDI	99I OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94 \text{ GeV}$

¹ ABBIENDI 99I give the 95%CL limit on the partial width $\Gamma(Z^0 \rightarrow p\mu) < 4.4 \text{ KeV}$ and we have transformed it into a branching ratio.

 Γ_{56}/Γ

NODE=S044PMU

NODE=S044PMU

NODE=S044PMU

NODE=S044PMU;LINKAGE=A

AVERAGE PARTICLE MULTIPLICITIES IN HADRONIC Z DECAY

Summed over particle and antiparticle, when appropriate.

NODE=S044260

NODE=S044260

 $\langle N_\gamma \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
20.97 ± 0.02 ± 1.15	ACKERSTAFF	98A OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$

NODE=S044GAM

NODE=S044GAM

 $\langle N_{\pi^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
17.03 ± 0.16 OUR AVERAGE			
17.007 ± 0.209	ABE	04C SLD	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$
17.26 ± 0.10 ± 0.88	ABREU	98L DLPH	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$
17.04 ± 0.31	BARATE	98V ALEP	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$
17.05 ± 0.43	AKERS	94P OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$

NODE=S044PIC

NODE=S044PIC

 $\langle N_{\pi^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
9.76 ± 0.26 OUR AVERAGE			
9.55 ± 0.06 ± 0.75	ACKERSTAFF	98A OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$
9.63 ± 0.13 ± 0.63	BARATE	97J ALEP	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$
9.90 ± 0.02 ± 0.33	ACCIARRI	96 L3	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$
9.2 ± 0.2 ± 1.0	ADAM	96 DLPH	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$

NODE=S044PIZ

NODE=S044PIZ


 $\langle N_\eta \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.01 ± 0.08 OUR AVERAGE			
1.20 ± 0.04 ± 0.11	HEISTER	02C ALEP	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$
0.97 ± 0.03 ± 0.11	ACKERSTAFF	98A OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$
0.93 ± 0.01 ± 0.09	ACCIARRI	96 L3	$E_{\text{cm}}^{\text{ee}} = 91.2 \text{ GeV}$

NODE=S044ETA

NODE=S044ETA

Error includes scale factor of 1.3. See the ideogram below.



$\langle N_{\rho^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
2.57 ± 0.15 OUR AVERAGE			
$2.59 \pm 0.03 \pm 0.16$	¹ BEDDALL 09		ALEPH archive, $E_{cm}^{ee} = 91.2$ GeV
$2.40 \pm 0.06 \pm 0.43$	ACKERSTAFF 98A	OPAL	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044RHC
 NODE=S044RHC

¹ BEDDALL 09 analyse 3.2 million hadronic Z decays as archived by ALEPH collaboration and report a value of $2.59 \pm 0.03 \pm 0.15 \pm 0.04$. The first error is statistical, the second systematic, and the third arises from extrapolation to full phase space. We combine the systematic errors in quadrature.

NODE=S044RHC;LINKAGE=BE

$\langle N_{\rho^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.24 ± 0.10 OUR AVERAGE	Error includes scale factor of 1.1.		
1.19 ± 0.10	ABREU 99J	DLPH	$E_{cm}^{ee} = 91.2$ GeV
$1.45 \pm 0.06 \pm 0.20$	BUSKULIC 96H	ALEP	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044RHZ
 NODE=S044RHZ

$\langle N_\omega \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.02 ± 0.06 OUR AVERAGE			
$1.00 \pm 0.03 \pm 0.06$	HEISTER 02C	ALEP	$E_{cm}^{ee} = 91.2$ GeV
$1.04 \pm 0.04 \pm 0.14$	ACKERSTAFF 98A	OPAL	$E_{cm}^{ee} = 91.2$ GeV
$1.17 \pm 0.09 \pm 0.15$	ACCIARRI 97D	L3	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044OME
 NODE=S044OME

$\langle N_{\eta'} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.05 OUR AVERAGE	Error includes scale factor of 2.4.		
$0.14 \pm 0.01 \pm 0.02$	ACKERSTAFF 98A	OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.25 ± 0.04	¹ ACCIARRI 97D	L3	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$0.068 \pm 0.018 \pm 0.016$	² BUSKULIC 92D	ALEP	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044ETP
 NODE=S044ETP

¹ ACCIARRI 97D obtain this value averaging over the two decay channels $\eta' \rightarrow \pi^+ \pi^- \eta$ and $\eta' \rightarrow \rho^0 \gamma$.

NODE=S044ETP;LINKAGE=B

² BUSKULIC 92D obtain this value for $x > 0.1$.

NODE=S044ETP;LINKAGE=A

$\langle N_{f_0(980)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.147 ± 0.011 OUR AVERAGE			
0.164 ± 0.021	ABREU 99J	DLPH	$E_{cm}^{ee} = 91.2$ GeV
$0.141 \pm 0.007 \pm 0.011$	ACKERSTAFF 98Q	OPAL	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044FZ
 NODE=S044FZ

$\langle N_{a_0(980)} \rangle$


VALUE	DOCUMENT ID	TECN	COMMENT
$0.27 \pm 0.04 \pm 0.10$	ACKERSTAFF 98A	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044AZC
NODE=S044AZC

 $\langle N_\phi \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.098 ± 0.006 OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.		
0.105 ± 0.008	ABE	99E SLD	$E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.091 \pm 0.002 \pm 0.003$	ACKERSTAFF	98Q OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.104 \pm 0.003 \pm 0.007$	ABREU	96U DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.122 \pm 0.004 \pm 0.008$	BUSKULIC	96H ALEP	$E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044PHI
NODE=S044PHI

 $\langle N_{f_2(1270)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.169 ± 0.025 OUR AVERAGE	Error includes scale factor of 1.4.		
0.214 ± 0.038	ABREU	99J DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.155 \pm 0.011 \pm 0.018$	ACKERSTAFF	98Q OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044F2
NODE=S044F2

 $\langle N_{f_1(1285)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.165 ± 0.051	¹ ABDALLAH	03H DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044F85
NODE=S044F85

¹ ABDALLAH 03H assume a $K\bar{K}\pi$ branching ratio of $(9.0 \pm 0.4)\%$.

NODE=S044F85;LINKAGE=A

 $\langle N_{f_1(1420)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.056 ± 0.012	¹ ABDALLAH	03H DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044F20
NODE=S044F20

¹ ABDALLAH 03H assume a $K\bar{K}\pi$ branching ratio of 100%.

NODE=S044F20;LINKAGE=A

 $\langle N_{f_2'(1525)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.012 ± 0.006	ABREU	99J DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044F2P
NODE=S044F2P

 $\langle N_{K^\pm} \rangle$


VALUE	DOCUMENT ID	TECN	COMMENT
2.24 ± 0.04 OUR AVERAGE			
2.203 ± 0.071	ABE	04C SLD	$E_{cm}^{ee} = 91.2 \text{ GeV}$
$2.21 \pm 0.05 \pm 0.05$	ABREU	98L DLPH	$E_{cm}^{ee} = 91.2 \text{ GeV}$
2.26 ± 0.12	BARATE	98v ALEP	$E_{cm}^{ee} = 91.2 \text{ GeV}$
2.42 ± 0.13	AKERS	94P OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044KC
NODE=S044KC

$\langle N_{K^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
2.039\pm0.025 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.		
2.093 \pm 0.004 \pm 0.029	BARATE	00O ALEP	$E_{cm}^{ee} = 91.2$ GeV
2.01 \pm 0.08	ABE	99E SLD	$E_{cm}^{ee} = 91.2$ GeV
2.024 \pm 0.006 \pm 0.042	ACCIARRI	97L L3	$E_{cm}^{ee} = 91.2$ GeV
1.962 \pm 0.022 \pm 0.056	ABREU	95L DLPH	$E_{cm}^{ee} = 91.2$ GeV
1.99 \pm 0.01 \pm 0.04	AKERS	95U OPAL	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044KZ
NODE=S044KZ

 $\langle N_{K^*(892)^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.72 \pm0.05 OUR AVERAGE			
0.712 \pm 0.031 \pm 0.059	ABREU	95L DLPH	$E_{cm}^{ee} = 91.2$ GeV
0.72 \pm 0.02 \pm 0.08	ACTON	93 OPAL	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044KSC
NODE=S044KSC

 $\langle N_{K^*(892)^0} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.739\pm0.022 OUR AVERAGE			
0.707 \pm 0.041	ABE	99E SLD	$E_{cm}^{ee} = 91.2$ GeV
0.74 \pm 0.02 \pm 0.02	ACKERSTAFF	97S OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.77 \pm 0.02 \pm 0.07	ABREU	96U DLPH	$E_{cm}^{ee} = 91.2$ GeV
0.83 \pm 0.01 \pm 0.09	BUSKULIC	96H ALEP	$E_{cm}^{ee} = 91.2$ GeV
0.97 \pm 0.18 \pm 0.31	ABREU	93 DLPH	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044KSZ
NODE=S044KSZ

 $\langle N_{K_2^*(1430)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.073\pm0.023	ABREU	99J DLPH	$E_{cm}^{ee} = 91.2$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.19 \pm 0.04 \pm 0.06	¹ AKERS	95X OPAL	$E_{cm}^{ee} = 91.2$ GeV

¹ AKERS 95X obtain this value for $x < 0.3$.

NODE=S044KS2
NODE=S044KS2

NODE=S044KS2;LINKAGE=A


 $\langle N_{D^\pm} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.187\pm0.020 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.		
0.170 \pm 0.009 \pm 0.014	ALEXANDER	96R OPAL	$E_{cm}^{ee} = 91.2$ GeV
0.251 \pm 0.026 \pm 0.025	BUSKULIC	94J ALEP	$E_{cm}^{ee} = 91.2$ GeV
0.199 \pm 0.019 \pm 0.024	¹ ABREU	93I DLPH	$E_{cm}^{ee} = 91.2$ GeV

NODE=S044DC
NODE=S044DC

¹ See ABREU 95 (erratum).

NODE=S044DC;LINKAGE=A

 **$\langle N_{D^0} \rangle$**

VALUE

0.462±0.026 OUR AVERAGE

0.465±0.017±0.027

0.518±0.052±0.035

0.403±0.038±0.044

DOCUMENT ID

TECN

COMMENT

ALEXANDER

96R

OPAL

 $E_{cm}^{ee} = 91.2$ GeV

BUSKULIC

94J

ALEP

 $E_{cm}^{ee} = 91.2$ GeV¹ ABREU

93I

DLPH

 $E_{cm}^{ee} = 91.2$ GeV¹ See ABREU 95 (erratum).NODE=S044DZ
NODE=S044DZ **$\langle N_{D_s^\pm} \rangle$**

VALUE

0.131±0.010±0.018

DOCUMENT ID

TECN

COMMENT

ALEXANDER

96R

OPAL

 $E_{cm}^{ee} = 91.2$ GeVNODE=S044DSP
NODE=S044DSP **$\langle N_{D^{*+}(2010)^+} \rangle$**

VALUE

0.183 ±0.008 OUR AVERAGE

0.1854±0.0041±0.0091

0.187 ±0.015 ±0.013

0.171 ±0.012 ±0.016

¹ ACKERSTAFF

98E

OPAL

 $E_{cm}^{ee} = 91.2$ GeV

BUSKULIC

94J

ALEP

 $E_{cm}^{ee} = 91.2$ GeV² ABREU

93I

DLPH

 $E_{cm}^{ee} = 91.2$ GeV

¹ ACKERSTAFF 98E systematic error includes an uncertainty of ±0.0069 due to the branching ratios $B(D^{*+} \rightarrow D^0 \pi^+) = 0.683 \pm 0.014$ and $B(D^0 \rightarrow K^- \pi^+) = 0.0383 \pm 0.0012$.

² See ABREU 95 (erratum).

NODE=S044DZ;LINKAGE=A

NODE=S044DSC
NODE=S044DSC

NODE=S044DSC;LINKAGE=B

NODE=S044DSC;LINKAGE=A

 $\langle N_{D_{s1}(2536)^+} \rangle$ VALUE (units 10^{-3})

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $2.9^{+0.7}_{-0.6} \pm 0.2$ ¹ ACKERSTAFF

97W

OPAL

 $E_{cm}^{ee} = 91.2$ GeV

¹ ACKERSTAFF 97W obtain this value for $x > 0.6$ and with the assumption that its decay width is saturated by the $D^* K$ final states.

NODE=S044DS1;LINKAGE=A

 $\langle N_{B^*} \rangle$

VALUE

0.28±0.01±0.03

DOCUMENT ID

TECN

COMMENT

¹ ABREU

95R

DLPH

 $E_{cm}^{ee} = 91.2$ GeVNODE=S044BS
NODE=S044BS¹ ABREU 95R quote this value for a flavor-averaged excited state.

NODE=S044BS;LINKAGE=A

$\langle N_{J/\psi(1S)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.0056 \pm 0.0003 \pm 0.0004$	¹ ALEXANDER 96B	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

¹ ALEXANDER 96B identify $J/\psi(1S)$ from the decays into lepton pairs.

NODE=S044JPS
NODE=S044JPS

NODE=S044JPS;LINKAGE=B

 $\langle N_{\psi(2S)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.0023 \pm 0.0004 \pm 0.0003$	ALEXANDER 96B	OPAL	$E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044P2S
NODE=S044P2S

 $\langle N_p \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
1.046 ± 0.026 OUR AVERAGE			
1.054 ± 0.035	ABE	04C	SLD $E_{cm}^{ee} = 91.2 \text{ GeV}$
$1.08 \pm 0.04 \pm 0.03$	ABREU	98L	DLPH $E_{cm}^{ee} = 91.2 \text{ GeV}$
1.00 ± 0.07	BARATE	98V	ALEP $E_{cm}^{ee} = 91.2 \text{ GeV}$
0.92 ± 0.11	AKERS	94P	OPAL $E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044PRO
NODE=S044PRO

 $\langle N_{\Delta(1232)++} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.087 ± 0.033 OUR AVERAGE	Error includes scale factor of 2.4.		
$0.079 \pm 0.009 \pm 0.011$	ABREU	95W	DLPH $E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.22 \pm 0.04 \pm 0.04$	ALEXANDER	95D	OPAL $E_{cm}^{ee} = 91.2 \text{ GeV}$


NODE=S044D12
NODE=S044D12

 $\langle N_\Lambda \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.388 ± 0.009 OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.		
$0.404 \pm 0.002 \pm 0.007$	BARATE	00O	ALEP $E_{cm}^{ee} = 91.2 \text{ GeV}$
0.395 ± 0.022	ABE	99E	SLD $E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.364 \pm 0.004 \pm 0.017$	ACCIARRI	97L	L3 $E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.374 \pm 0.002 \pm 0.010$	ALEXANDER	97D	OPAL $E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.357 \pm 0.003 \pm 0.017$	ABREU	93L	DLPH $E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044LAM
NODE=S044LAM

WEIGHTED AVERAGE
 0.388 ± 0.009 (Error scaled by 1.7)

 $\langle N_\Lambda \rangle$ $\langle N_{\Lambda(1520)} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0224 ± 0.0027 OUR AVERAGE			
$0.029 \pm 0.005 \pm 0.005$	ABREU	00P	DLPH $E_{cm}^{ee} = 91.2 \text{ GeV}$
$0.0213 \pm 0.0021 \pm 0.0019$	ALEXANDER	97D	OPAL $E_{cm}^{ee} = 91.2 \text{ GeV}$

NODE=S044L15
NODE=S044L15

$\langle N_{\Sigma^+} \rangle$

VALUE

0.107 ± 0.010 OUR AVERAGE

0.114 ± 0.011 ± 0.009

0.099 ± 0.008 ± 0.013

DOCUMENT ID

TECN

COMMENT

ACCIARRI

00J

L3

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ALEXANDER

97E

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044SI+
NODE=S044SI+ $\langle N_{\Sigma^-} \rangle$

VALUE

0.082 ± 0.007 OUR AVERAGE

0.081 ± 0.002 ± 0.010

0.083 ± 0.006 ± 0.009

DOCUMENT ID

TECN

COMMENT

ABREU

00P

DLPH

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ALEXANDER

97E

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044SI-
NODE=S044SI- $\langle N_{\Sigma^+ + \Sigma^-} \rangle$

VALUE

0.181 ± 0.018 OUR AVERAGE

0.182 ± 0.010 ± 0.016

0.170 ± 0.014 ± 0.061

¹

ALEXANDER

97E

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ABREU

950

DLPH

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044SIC
NODE=S044SIC

¹We have combined the values of $\langle N_{\Sigma^+} \rangle$ and $\langle N_{\Sigma^-} \rangle$ from ALEXANDER 97E adding the statistical and systematic errors of the two final states separately in quadrature. If isospin symmetry is assumed this value becomes $0.174 \pm 0.010 \pm 0.015$.

NODE=S044SIC;LINKAGE=A

 $\langle N_{\Sigma^0} \rangle$

VALUE

0.076 ± 0.010 OUR AVERAGE

0.095 ± 0.015 ± 0.013

0.071 ± 0.012 ± 0.013

0.070 ± 0.010 ± 0.010

DOCUMENT ID

TECN

COMMENT

ACCIARRI

00J

L3

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ALEXANDER

97E

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ADAM

96B

DLPH

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044SIZ
NODE=S044SIZ $\langle N_{(\Sigma^+ + \Sigma^- + \Sigma^0)/3} \rangle$

VALUE

0.084 ± 0.005 ± 0.008

DOCUMENT ID

TECN

COMMENT

ALEXANDER

97E

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044SIG
NODE=S044SIG $\langle N_{\Sigma(1385)^+} \rangle$

VALUE

0.0239 ± 0.0009 ± 0.0012

DOCUMENT ID

TECN

COMMENT

ALEXANDER

97D

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S04413+
NODE=S04413+ $\langle N_{\Sigma(1385)^-} \rangle$

VALUE

0.0240 ± 0.0010 ± 0.0014

DOCUMENT ID

TECN

COMMENT

ALEXANDER

97D

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S04413-
NODE=S04413- $\langle N_{\Sigma(1385)^+ + \Sigma(1385)^-} \rangle$

VALUE

0.046 ± 0.004 OUR AVERAGE

0.0479 ± 0.0013 ± 0.0026

0.0382 ± 0.0028 ± 0.0045

Error includes scale factor of 1.6.

ALEXANDER

97D

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ABREU

950

DLPH

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044S13
NODE=S044S13 $\langle N_{\Xi^-} \rangle$

VALUE

0.0258 ± 0.0009 OUR AVERAGE

0.0247 ± 0.0009 ± 0.0025

0.0259 ± 0.0004 ± 0.0009

DOCUMENT ID

TECN

COMMENT

ABDALLAH

06E

DLPH

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ALEXANDER

97D

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044XI-
NODE=S044XI- $\langle N_{\Xi(1530)^0} \rangle$

VALUE

0.0059 ± 0.0011 OUR AVERAGE

0.0045 ± 0.0005 ± 0.0006

0.0068 ± 0.0005 ± 0.0004

Error includes scale factor of 2.3.

ABDALLAH

05C

DLPH

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ALEXANDER

97D

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044X15
NODE=S044X15 $\langle N_{\Omega^-} \rangle$

VALUE

0.00164 ± 0.00028 OUR AVERAGE

0.0018 ± 0.0003 ± 0.0002

0.0014 ± 0.0002 ± 0.0004

DOCUMENT ID

TECN

COMMENT

ALEXANDER

97D

OPAL

 $E_{cm}^{ee} = 91.2 \text{ GeV}$

ADAM

96B

DLPH

 $E_{cm}^{ee} = 91.2 \text{ GeV}$ NODE=S044OM-
NODE=S044OM-

$\langle N_{\Lambda_c^+} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
0.078±0.012±0.012	ALEXANDER	96R	OPAL $E_{cm}^{ee} = 91.2$ GeV

NODE=S044L+C
 NODE=S044L+C

 $\langle N_D \rangle$

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

NODE=S044DBR
 NODE=S044DBR

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.9±1.8±0.5 ¹SCHAELE 06A ALEP $E_{cm}^{ee} = 91.2$ GeV

¹SCHAELE 06A obtain this anti-deuteron production rate per hadronic Z decay in the anti-deuteron momentum range from 0.62 to 1.03 GeV/c.


NODE=S044DBR;LINKAGE=SC

 $\langle N_{charged} \rangle$

VALUE	DOCUMENT ID	TECN	COMMENT
20.76±0.16 OUR AVERAGE	Error includes scale factor of 2.1.	See the ideogram below.	
20.46±0.01±0.11	ACHARD	03G	L3 $E_{cm}^{ee} = 91.2$ GeV
21.21±0.01±0.20	ABREU	99	DLPH $E_{cm}^{ee} = 91.2$ GeV
21.05±0.20	AKERS	95Z	OPAL $E_{cm}^{ee} = 91.2$ GeV
20.91±0.03±0.22	BUSKULIC	95R	ALEP $E_{cm}^{ee} = 91.2$ GeV
21.40±0.43	ACTON	92B	OPAL $E_{cm}^{ee} = 91.2$ GeV
20.71±0.04±0.77	ABREU	91H	DLPH $E_{cm}^{ee} = 91.2$ GeV
20.7 ±0.7	ADEVA	91I	L3 $E_{cm}^{ee} = 91.2$ GeV
20.1 ±1.0 ±0.9	ABRAMS	90	MRK2 $E_{cm}^{ee} = 91.1$ GeV

NODE=S044CHG
 NODE=S044CHG

WEIGHTED AVERAGE
 20.76±0.16 (Error scaled by 2.1)



Z HADRONIC POLE CROSS SECTION

NODE=S044SH

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note "The Z boson" and ref. LEP-SLC 06). This quantity is defined as

$$\sigma_h^0 = \frac{12\pi}{M_Z^2} \frac{\Gamma(e^+e^-)\Gamma(\text{hadrons})}{\Gamma_Z^2}$$

It is one of the parameters used in the Z lineshape fit.

NODE=S044SH

VALUE (nb)	EVTs	DOCUMENT ID	TECN	COMMENT
41.541±0.037 OUR FIT				
41.501±0.055	4.10M	¹ ABBIENDI	01A	OPAL $E_{cm}^{ee} = 88-94$ GeV
41.578±0.069	3.70M	ABREU	00F	DLPH $E_{cm}^{ee} = 88-94$ GeV
41.535±0.055	3.54M	ACCIARRI	00C	L3 $E_{cm}^{ee} = 88-94$ GeV
41.559±0.058	4.07M	² BARATE	00C	ALEP $E_{cm}^{ee} = 88-94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
42 ±4	450	ABRAMS	89B	MRK2 $E_{cm}^{ee} = 89.2-93.0$ GeV

NODE=S044SH

¹ ABBIENDI 01A error includes approximately 0.031 due to statistics, 0.033 due to event selection systematics, 0.029 due to uncertainty in luminosity measurement, and 0.011 due to LEP energy uncertainty.

² BARATE 00C error includes approximately 0.030 due to statistics, 0.026 due to experimental systematics, and 0.025 due to uncertainty in luminosity measurement.

Z VECTOR COUPLINGS

These quantities are the effective vector couplings of the Z to charged leptons. Their magnitude is derived from a measurement of the Z lineshape and the forward-backward lepton asymmetries as a function of energy around the Z mass. The relative sign among the vector to axial-vector couplings is obtained from a measurement of the Z asymmetry parameters, A_e , A_μ , and A_τ . By convention the sign of g_A^e is fixed to be negative (and opposite to that of g_V^e obtained using ν_e scattering measurements). For the light quarks, the sign of the couplings is assigned consistently with this assumption. The fit values quoted below correspond to global nine- or five-parameter fits to lineshape, lepton forward-backward asymmetry, and A_e , A_μ , and A_τ measurements. See the note "The Z boson" and ref. LEP-SLC 06 for details. Where $p\bar{p}$ and $e p$ data is quoted, OUR FIT value corresponds to a weighted average of this with the LEP/SLD fit result.

g_V^e

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
−0.03817±0.00047 OUR FIT				
−0.058 ±0.016 ±0.007	5026	¹ ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV
−0.0346 ±0.0023	137.0K	² ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
−0.0412 ±0.0027	124.4k	³ ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
−0.0400 ±0.0037		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
−0.0414 ±0.0020		⁴ ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV

¹ ACOSTA 05M determine the forward-backward asymmetry of e^+e^- pairs produced via $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+e^-$ in 15 M(e^+e^-) effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the Z to e^+e^- , assuming the quark couplings are as predicted by the standard model. Higher order radiative corrections have not been taken into account.

² ABBIENDI 010 use their measurement of the τ polarization in addition to the lineshape and forward-backward lepton asymmetries.

³ ACCIARRI 00C use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

⁴ ABE 95J obtain this result combining polarized Bhabha results with the A_{LR} measurement of ABE 94C. The Bhabha results alone give $-0.0507 \pm 0.0096 \pm 0.0020$.

g_V^μ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
−0.0367±0.0023 OUR FIT				
−0.0388 ^{+0.0060} _{−0.0064}	182.8K	¹ ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
−0.0386±0.0073	113.4k	² ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
−0.0362±0.0061		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
−0.0413±0.0060	66143	³ ABBIENDI	01K OPAL	$E_{cm}^{ee} = 89-93$ GeV

¹ ABBIENDI 010 use their measurement of the τ polarization in addition to the lineshape and forward-backward lepton asymmetries.

² ACCIARRI 00C use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

³ ABBIENDI 01K obtain this from an angular analysis of the muon pair asymmetry which takes into account effects of initial state radiation on an event by event basis and of initial-final state interference.

g_V^τ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
−0.0366±0.0010 OUR FIT				
−0.0365±0.0023	151.5K	¹ ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
−0.0384±0.0026	103.0k	² ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
−0.0361±0.0068		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

¹ ABBIENDI 010 use their measurement of the τ polarization in addition to the lineshape and forward-backward lepton asymmetries.

² ACCIARRI 00C use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

NODE=S044SH;LINKAGE=DB

NODE=S044SH;LINKAGE=AC

NODE=S044223

NODE=S044223

NODE=S044GEV
NODE=S044GEV

NODE=S044GEV;LINKAGE=AC

NODE=S044GEV;LINKAGE=OA

NODE=S044GEV;LINKAGE=Z

NODE=S044GEV;LINKAGE=KG

NODE=S044GMV
NODE=S044GMV

NODE=S044GMV;LINKAGE=OA

NODE=S044GMV;LINKAGE=Z

NODE=S044GMV;LINKAGE=GM

NODE=S044GTV
NODE=S044GTV

NODE=S044GTV;LINKAGE=OA

NODE=S044GTV;LINKAGE=Z

g_V^ℓ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.03783 ± 0.00041 OUR FIT				
-0.0358 ± 0.0014	471.3K	¹ ABBIENDI	01O OPAL	$E_{cm}^{ee} = 88-94$ GeV
-0.0397 ± 0.0020	379.4k	² ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
-0.0397 ± 0.0017	340.8k	³ ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
-0.0383 ± 0.0018	500k	BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

¹ ABBIENDI 01O use their measurement of the τ polarization in addition to the lineshape and forward-backward lepton asymmetries.

² Using forward-backward lepton asymmetries.

³ ACCIARRI 00C use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

NODE=S044GLV
NODE=S044GLV

NODE=S044GLV;LINKAGE=OA

NODE=S044GLV;LINKAGE=B

NODE=S044GLV;LINKAGE=Z

 g_V^u

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.25 ± 0.07 -0.06 OUR AVERAGE				
0.201 ± 0.112	156k	¹ ABAZOV	11D D0	$E_{cm}^{p\bar{p}} = 1.97$ TeV
0.27 ± 0.13	1500	² AKTAS	06 H1	$e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X$, $\sqrt{s} \approx 300$ GeV
0.24 ± 0.28 -0.11		³ LEP-SLC	06	$E_{cm}^{ee} = 88-94$ GeV
0.399 ± 0.152 -0.188 ± 0.066	5026	⁴ ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV

¹ ABAZOV 11D study $p\bar{p} \rightarrow Z/\gamma^* e^+ e^-$ events using 5 fb^{-1} data at $\sqrt{s} = 1.96$ TeV. The candidate events are selected by requiring two isolated electromagnetic showers with $E_T > 25$ GeV, at least one electron in the central region and the di-electron mass in the range 50–1000 GeV. From the forward-backward asymmetry, determined as a function of the di-electron mass, they derive the axial and vector couplings of the u - and d - quarks and the value of $\sin^2 \theta_{eff}^\ell = 0.2309 \pm 0.0008(\text{stat}) \pm 0.0006(\text{syst})$.

² AKTAS 06 fit the neutral current ($1.5 \leq Q^2 \leq 30,000 \text{ GeV}^2$) and charged current ($1.5 \leq Q^2 \leq 15,000 \text{ GeV}^2$) differential cross sections. In the determination of the u -quark couplings the electron and d -quark couplings are fixed to their standard model values.

³ LEP-SLC 06 is a combination of the results from LEP and SLC experiments using light quark tagging. s - and d -quark couplings are assumed to be identical.

⁴ ACOSTA 05M determine the forward-backward asymmetry of $e^+ e^-$ pairs produced via $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+ e^-$ in 15 $M(e^+ e^-)$ effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the Z to the light quarks, assuming the electron couplings are as predicted by the Standard Model. Higher order radiative corrections have not been taken into account.

NODE=S044GUV
NODE=S044GUV

NODE=S044GUV;LINKAGE=AB

NODE=S044GUV;LINKAGE=AK

NODE=S044GUV;LINKAGE=LS

NODE=S044GUV;LINKAGE=AC

 g_V^d

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.33 ± 0.05 -0.06 OUR AVERAGE				
-0.351 ± 0.251	156k	¹ ABAZOV	11D D0	$E_{cm}^{p\bar{p}} = 1.97$ TeV
-0.33 ± 0.33	1500	² AKTAS	06 H1	$e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X$, $\sqrt{s} \approx 300$ GeV
-0.33 ± 0.05 -0.07		³ LEP-SLC	06	$E_{cm}^{ee} = 88-94$ GeV
-0.226 ± 0.635 -0.290 ± 0.090	5026	⁴ ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV

¹ ABAZOV 11D study $p\bar{p} \rightarrow Z/\gamma^* e^+ e^-$ events using 5 fb^{-1} data at $\sqrt{s} = 1.96$ TeV. The candidate events are selected by requiring two isolated electromagnetic showers with $E_T > 25$ GeV, at least one electron in the central region and the di-electron mass in the range 50–1000 GeV. From the forward-backward asymmetry, determined as a function of the di-electron mass, they derive the axial and vector couplings of the u - and d - quarks and the value of $\sin^2 \theta_{eff}^\ell = 0.2309 \pm 0.0008(\text{stat}) \pm 0.0006(\text{syst})$.

² AKTAS 06 fit the neutral current ($1.5 \leq Q^2 \leq 30,000 \text{ GeV}^2$) and charged current ($1.5 \leq Q^2 \leq 15,000 \text{ GeV}^2$) differential cross sections. In the determination of the d -quark couplings the electron and u -quark couplings are fixed to their standard model values.

³ LEP-SLC 06 is a combination of the results from LEP and SLC experiments using light quark tagging. s - and d -quark couplings are assumed to be identical.

⁴ ACOSTA 05M determine the forward-backward asymmetry of $e^+ e^-$ pairs produced via $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+ e^-$ in 15 $M(e^+ e^-)$ effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the Z to the light quarks, assuming the electron couplings are as predicted by the Standard Model. Higher order radiative corrections have not been taken into account.

NODE=S044GDV
NODE=S044GDV

NODE=S044GDV;LINKAGE=AB

NODE=S044GDV;LINKAGE=AK

NODE=S044GDV;LINKAGE=LS

NODE=S044GDV;LINKAGE=AC

Z AXIAL-VECTOR COUPLINGS

These quantities are the effective axial-vector couplings of the Z to charged leptons. Their magnitude is derived from a measurement of the Z lineshape and the forward-backward lepton asymmetries as a function of energy around the Z mass. The relative sign among the vector to axial-vector couplings is obtained from a measurement of the Z asymmetry parameters, A_e , A_μ , and A_τ . By convention the sign of g_A^e is fixed to be negative (and opposite to that of g_V^e obtained using ν_e scattering measurements). For the light quarks, the sign of the couplings is assigned consistently with this assumption. The fit values quoted below correspond to global nine- or five-parameter fits to lineshape, lepton forward-backward asymmetry, and A_e , A_μ , and A_τ measurements. See the note "The Z boson" and ref. LEP-SLC 06 for details. Where $p\bar{p}$ and $e p$ data is quoted, OUR FIT value corresponds to a weighted average of this with the LEP/SLD fit result.

g_A^e

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
−0.50111 ± 0.00035 OUR FIT				
−0.528 ± 0.123 ± 0.059	5026	¹ ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV
−0.50062 ± 0.00062	137.0K	² ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
−0.5015 ± 0.0007	124.4k	³ ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
−0.50166 ± 0.00057		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
−0.4977 ± 0.0045		⁴ ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV

¹ ACOSTA 05M determine the forward-backward asymmetry of e^+e^- pairs produced via $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+e^-$ in 15 $M(e^+e^-)$ effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the Z to e^+e^- , assuming the quark couplings are as predicted by the standard model. Higher order radiative corrections have not been taken into account.

² ABBIENDI 010 use their measurement of the τ polarization in addition to the lineshape and forward-backward lepton asymmetries.

³ ACCIARRI 00C use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

⁴ ABE 95J obtain this result combining polarized Bhabha results with the A_{LR} measurement of ABE 94C. The Bhabha results alone give $-0.4968 \pm 0.0039 \pm 0.0027$.

NODE=S044GEA
NODE=S044GEA

NODE=S044GEA;LINKAGE=AC

NODE=S044GEA;LINKAGE=OA

NODE=S044GEA;LINKAGE=Z

NODE=S044GEA;LINKAGE=KG

g_A^μ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
−0.50120 ± 0.00054 OUR FIT				
−0.50117 ± 0.00099	182.8K	¹ ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
−0.5009 ± 0.0014	113.4k	² ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
−0.50046 ± 0.00093		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
−0.520 ± 0.015	66143	³ ABBIENDI	01K OPAL	$E_{cm}^{ee} = 89-93$ GeV

¹ ABBIENDI 010 use their measurement of the τ polarization in addition to the lineshape and forward-backward lepton asymmetries.

² ACCIARRI 00C use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

³ ABBIENDI 01K obtain this from an angular analysis of the muon pair asymmetry which takes into account effects of initial state radiation on an event by event basis and of initial-final state interference.

NODE=S044GMA
NODE=S044GMA

NODE=S044GMA;LINKAGE=OA

NODE=S044GMA;LINKAGE=Z

NODE=S044GMA;LINKAGE=GM

g_A^τ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
−0.50204 ± 0.00064 OUR FIT				
−0.50165 ± 0.00124	151.5K	¹ ABBIENDI	010 OPAL	$E_{cm}^{ee} = 88-94$ GeV
−0.5023 ± 0.0017	103.0k	² ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
−0.50216 ± 0.00100		BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

¹ ABBIENDI 010 use their measurement of the τ polarization in addition to the lineshape and forward-backward lepton asymmetries.

² ACCIARRI 00C use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

NODE=S044GTA
NODE=S044GTA

NODE=S044GTA;LINKAGE=OA

NODE=S044GTA;LINKAGE=Z

g_A^ℓ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.50123 ± 0.00026 OUR FIT				
-0.50089 ± 0.00045	471.3K	¹ ABBIENDI	01O OPAL	$E_{cm}^{ee} = 88-94$ GeV
-0.5007 ± 0.0005	379.4k	ABREU	00F DLPH	$E_{cm}^{ee} = 88-94$ GeV
-0.50153 ± 0.00053	340.8k	² ACCIARRI	00C L3	$E_{cm}^{ee} = 88-94$ GeV
-0.50150 ± 0.00046	500k	BARATE	00C ALEP	$E_{cm}^{ee} = 88-94$ GeV

¹ ABBIENDI 01O use their measurement of the τ polarization in addition to the lineshape and forward-backward lepton asymmetries.

² ACCIARRI 00C use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

NODE=S044GLA
NODE=S044GLA

NODE=S044GLA;LINKAGE=OA

NODE=S044GLA;LINKAGE=Z

 g_A^u

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.50^{+0.04}_{-0.06}$ OUR AVERAGE				
0.501 ± 0.110	156k	¹ ABAZOV	11D D0	$E_{cm}^{p\bar{p}} = 1.97$ TeV
0.57 ± 0.08	1500	² AKTAS	06 H1	$e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X$, $\sqrt{s} \approx 300$ GeV
$0.47^{+0.05}_{-0.33}$		³ LEP-SLC	06	$E_{cm}^{ee} = 88-94$ GeV
$0.441^{+0.207}_{-0.173} \pm 0.067$	5026	⁴ ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV

¹ ABAZOV 11D study $p\bar{p} \rightarrow Z/\gamma^* e^+ e^-$ events using 5 fb^{-1} data at $\sqrt{s} = 1.96$ TeV. The candidate events are selected by requiring two isolated electromagnetic showers with $E_T > 25$ GeV, at least one electron in the central region and the di-electron mass in the range 50–1000 GeV. From the forward-backward asymmetry, determined as a function of the di-electron mass, they derive the axial and vector couplings of the u - and d - quarks and the value of $\sin^2 \theta_{eff}^\ell = 0.2309 \pm 0.0008(\text{stat}) \pm 0.0006(\text{syst})$.

² AKTAS 06 fit the neutral current ($1.5 \leq Q^2 \leq 30,000 \text{ GeV}^2$) and charged current ($1.5 \leq Q^2 \leq 15,000 \text{ GeV}^2$) differential cross sections. In the determination of the u -quark couplings the electron and d -quark couplings are fixed to their standard model values.

³ LEP-SLC 06 is a combination of the results from LEP and SLC experiments using light quark tagging. s - and d -quark couplings are assumed to be identical.

⁴ ACOSTA 05M determine the forward-backward asymmetry of $e^+ e^-$ pairs produced via $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+ e^-$ in $15 \text{ M}(e^+ e^-)$ effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the Z to the light quarks, assuming the electron couplings are as predicted by the Standard Model. Higher order radiative corrections have not been taken into account.

NODE=S044GUA
NODE=S044GUA

NODE=S044GUA;LINKAGE=AB

NODE=S044GUA;LINKAGE=AK

NODE=S044GUA;LINKAGE=LS

NODE=S044GUA;LINKAGE=AC

 g_A^d

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.523^{+0.050}_{-0.029}$ OUR AVERAGE				
-0.497 ± 0.165	156k	¹ ABAZOV	11D D0	$E_{cm}^{p\bar{p}} = 1.97$ TeV
-0.80 ± 0.24	1500	² AKTAS	06 H1	$e^\pm p \rightarrow \bar{\nu}_e(\nu_e)X$, $\sqrt{s} \approx 300$ GeV
$-0.52^{+0.05}_{-0.03}$		³ LEP-SLC	06	$E_{cm}^{ee} = 88-94$ GeV
$-0.016^{+0.346}_{-0.536} \pm 0.091$	5026	⁴ ACOSTA	05M CDF	$E_{cm}^{p\bar{p}} = 1.96$ TeV

¹ ABAZOV 11D study $p\bar{p} \rightarrow Z/\gamma^* e^+ e^-$ events using 5 fb^{-1} data at $\sqrt{s} = 1.96$ TeV. The candidate events are selected by requiring two isolated electromagnetic showers with $E_T > 25$ GeV, at least one electron in the central region and the di-electron mass in the range 50–1000 GeV. From the forward-backward asymmetry, determined as a function of the di-electron mass, they derive the axial and vector couplings of the u - and d - quarks and the value of $\sin^2 \theta_{eff}^\ell = 0.2309 \pm 0.0008(\text{stat}) \pm 0.0006(\text{syst})$.

² AKTAS 06 fit the neutral current ($1.5 \leq Q^2 \leq 30,000 \text{ GeV}^2$) and charged current ($1.5 \leq Q^2 \leq 15,000 \text{ GeV}^2$) differential cross sections. In the determination of the d -quark couplings the electron and u -quark couplings are fixed to their standard model values.

³ LEP-SLC 06 is a combination of the results from LEP and SLC experiments using light quark tagging. s - and d -quark couplings are assumed to be identical.

⁴ ACOSTA 05M determine the forward-backward asymmetry of $e^+ e^-$ pairs produced via $q\bar{q} \rightarrow Z/\gamma^* \rightarrow e^+ e^-$ in $15 \text{ M}(e^+ e^-)$ effective mass bins ranging from 40 GeV to 600 GeV. These results are used to obtain the vector and axial-vector couplings of the Z to the light quarks, assuming the electron couplings are as predicted by the Standard Model. Higher order radiative corrections have not been taken into account.

NODE=S044GDA
NODE=S044GDA

NODE=S044GDA;LINKAGE=AB

NODE=S044GDA;LINKAGE=AK

NODE=S044GDA;LINKAGE=LS

NODE=S044GDA;LINKAGE=AC

Z COUPLINGS TO NEUTRAL LEPTONS

Averaging over neutrino species, the invisible Z decay width determines the effective neutrino coupling $g^{\nu\ell}$. For $g^{\nu e}$ and $g^{\nu\mu}$, $\nu_e e$ and $\nu_\mu e$ scattering results are combined with g_A^e and g_V^e measurements at the Z mass to obtain $g^{\nu e}$ and $g^{\nu\mu}$ following NOVIKOV 93C.

$g^{\nu\ell}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.50076 ± 0.00076	¹ LEP-SLC 06		$E_{cm}^{ee} = 88-94$ GeV

¹ From invisible Z -decay width.

$g^{\nu e}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.528 ± 0.085	¹ VILAIN 94	CHM2	From $\nu_\mu e$ and $\nu_e e$ scattering

¹ VILAIN 94 derive this value from their value of $g^{\nu\mu}$ and their ratio $g^{\nu e}/g^{\nu\mu} = 1.05^{+0.15}_{-0.18}$.

$g^{\nu\mu}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.502 ± 0.017	¹ VILAIN 94	CHM2	From $\nu_\mu e$ scattering

¹ VILAIN 94 derive this value from their measurement of the couplings $g_A^{e\nu\mu} = -0.503 \pm 0.017$ and $g_V^{e\nu\mu} = -0.035 \pm 0.017$ obtained from $\nu_\mu e$ scattering. We have re-evaluated this value using the current PDG values for g_A^e and g_V^e .

Z ASYMMETRY PARAMETERS

For each fermion-antifermion pair coupling to the Z these quantities are defined as

$$A_f = \frac{2g_V^f g_A^f}{(g_V^f)^2 + (g_A^f)^2}$$

where g_V^f and g_A^f are the effective vector and axial-vector couplings. For their relation to the various lepton asymmetries see the note "The Z boson" and ref. LEP-SLC 06.

A_e

Using polarized beams, this quantity can also be measured as $(\sigma_L - \sigma_R)/(\sigma_L + \sigma_R)$, where σ_L and σ_R are the e^+e^- production cross sections for Z bosons produced with left-handed and right-handed electrons respectively.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1515 ± 0.0019 OUR AVERAGE				
0.1454 ± 0.0108 ± 0.0036	144810	¹ ABBIENDI	01O OPAL	$E_{cm}^{ee} = 88-94$ GeV
0.1516 ± 0.0021	559000	² ABE	01B SLD	$E_{cm}^{ee} = 91.24$ GeV
0.1504 ± 0.0068 ± 0.0008		³ HEISTER	01 ALEP	$E_{cm}^{ee} = 88-94$ GeV
0.1382 ± 0.0116 ± 0.0005	105000	⁴ ABREU	00E DLPH	$E_{cm}^{ee} = 88-94$ GeV
0.1678 ± 0.0127 ± 0.0030	137092	⁵ ACCIARRI	98H L3	$E_{cm}^{ee} = 88-94$ GeV
0.162 ± 0.041 ± 0.014	89838	⁶ ABE	97 SLD	$E_{cm}^{ee} = 91.27$ GeV
0.202 ± 0.038 ± 0.008		⁷ ABE	95J SLD	$E_{cm}^{ee} = 91.31$ GeV

¹ ABBIENDI 01O fit for A_e and A_τ from measurements of the τ polarization at varying τ production angles. The correlation between A_e and A_τ is less than 0.03.

² ABE 01B use the left-right production and left-right forward-backward decay asymmetries in leptonic Z decays to obtain a value of 0.1544 ± 0.0060 . This is combined with left-right production asymmetry measurement using hadronic Z decays (ABE 00B) to obtain the quoted value.

³ HEISTER 01 obtain this result fitting the τ polarization as a function of the polar production angle of the τ .

⁴ ABREU 00E obtain this result fitting the τ polarization as a function of the polar τ production angle. This measurement is a combination of different analyses (exclusive τ decay modes, inclusive hadronic 1-prong reconstruction, and a neural network analysis).

⁵ Derived from the measurement of forward-backward τ polarization asymmetry.

⁶ ABE 97 obtain this result from a measurement of the observed left-right charge asymmetry, $A_Q^{\text{obs}} = 0.225 \pm 0.056 \pm 0.019$, in hadronic Z decays. If they combine this value of A_Q^{obs} with their earlier measurement of A_{LR}^{obs} they determine A_e to be $0.1574 \pm 0.0197 \pm 0.0067$ independent of the beam polarization.

⁷ ABE 95J obtain this result from polarized Bhabha scattering.

NODE=S044228

NODE=S044228

NODE=S044GNL
NODE=S044GNL

NODE=S044GNL;LINKAGE=LE

NODE=S044GNE
NODE=S044GNE

NODE=S044GNE;LINKAGE=A

NODE=S044GNM
NODE=S044GNM

NODE=S044GNM;LINKAGE=A

NODE=S044224

NODE=S044224

NODE=S044AE
NODE=S044AE

NODE=S044AE

NODE=S044AE;LINKAGE=OA

NODE=S044AE;LINKAGE=BB

NODE=S044AE;LINKAGE=HE

NODE=S044AE;LINKAGE=L

NODE=S044AE;LINKAGE=A
NODE=S044AE;LINKAGE=E

NODE=S044AE;LINKAGE=KG

A_μ

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in $\mu^+\mu^-$ production at SLC using a polarized electron beam. This double asymmetry eliminates the dependence on the Z - e - e coupling parameter A_e .

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.142 ± 0.015	16844	¹ ABE	01B SLD	$E_{cm}^{ee} = 91.24$ GeV

¹ ABE 01B obtain this direct measurement using the left-right production and left-right forward-backward polar angle asymmetries in $\mu^+\mu^-$ decays of the Z boson obtained with a polarized electron beam.

NODE=S044AM

NODE=S044AM

NODE=S044AM

NODE=S044AM;LINKAGE=BB

 A_τ

The LEP Collaborations derive this quantity from the measurement of the τ polarization in $Z \rightarrow \tau^+\tau^-$. The SLD Collaboration directly extracts this quantity from its measured left-right forward-backward asymmetry in $Z \rightarrow \tau^+\tau^-$ produced using a polarized e^- beam. This double asymmetry eliminates the dependence on the Z - e - e coupling parameter A_e .

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.143 ± 0.004 OUR AVERAGE				
0.1456 ± 0.0076 ± 0.0057	144810	¹ ABBIENDI	01O OPAL	$E_{cm}^{ee} = 88-94$ GeV
0.136 ± 0.015	16083	² ABE	01B SLD	$E_{cm}^{ee} = 91.24$ GeV
0.1451 ± 0.0052 ± 0.0029		³ HEISTER	01 ALEP	$E_{cm}^{ee} = 88-94$ GeV
0.1359 ± 0.0079 ± 0.0055	105000	⁴ ABREU	00E DLPH	$E_{cm}^{ee} = 88-94$ GeV
0.1476 ± 0.0088 ± 0.0062	137092	ACCIARRI	98H L3	$E_{cm}^{ee} = 88-94$ GeV

¹ ABBIENDI 01O fit for A_e and A_τ from measurements of the τ polarization at varying τ production angles. The correlation between A_e and A_τ is less than 0.03.

² ABE 01B obtain this direct measurement using the left-right production and left-right forward-backward polar angle asymmetries in $\tau^+\tau^-$ decays of the Z boson obtained with a polarized electron beam.

³ HEISTER 01 obtain this result fitting the τ polarization as a function of the polar production angle of the τ .

⁴ ABREU 00E obtain this result fitting the τ polarization as a function of the polar τ production angle. This measurement is a combination of different analyses (exclusive τ decay modes, inclusive hadronic 1-prong reconstruction, and a neural network analysis).

NODE=S044AT

NODE=S044AT

NODE=S044AT

NODE=S044AT;LINKAGE=OA

NODE=S044AT;LINKAGE=BB

NODE=S044AT;LINKAGE=HE

NODE=S044AT;LINKAGE=L

 A_s

The SLD Collaboration directly extracts this quantity by a simultaneous fit to four measured s -quark polar angle distributions corresponding to two states of e^- polarization (positive and negative) and to the K^+K^- and $K^\pm K_S^0$ strange particle tagging modes in the hadronic final states.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.895 ± 0.066 ± 0.062	2870	¹ ABE	00D SLD	$E_{cm}^{ee} = 91.2$ GeV

¹ ABE 00D tag $Z \rightarrow s\bar{s}$ events by an absence of B or D hadrons and the presence in each hemisphere of a high momentum K^\pm or K_S^0 .

NODE=S044AS

NODE=S044AS

NODE=S044AS

NODE=S044AS;LINKAGE=A

 A_c

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in $c\bar{c}$ production at SLC using polarized electron beam. This double asymmetry eliminates the dependence on the Z - e - e coupling parameter A_e . OUR FIT is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the note "The Z boson" and ref. LEP-SLC 06.

VALUE	DOCUMENT ID	TECN	COMMENT
0.670 ± 0.027 OUR FIT			
0.6712 ± 0.0224 ± 0.0157	¹ ABE	05 SLD	$E_{cm}^{ee} = 91.24$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.583 ± 0.055 ± 0.055	² ABE	02G SLD	$E_{cm}^{ee} = 91.24$ GeV
0.688 ± 0.041	³ ABE	01C SLD	$E_{cm}^{ee} = 91.25$ GeV

¹ ABE 05 use hadronic Z decays collected during 1996–98 to obtain an enriched sample of $c\bar{c}$ events tagging on the invariant mass of reconstructed secondary decay vertices. The charge of the underlying c -quark is obtained with an algorithm that takes into account the net charge of the vertex as well as the charge of tracks emanating from the vertex and identified as kaons. This yields (9970 events) $A_c = 0.6747 \pm 0.0290 \pm 0.0233$. Taking into account all correlations with earlier results reported in ABE 02G and ABE 01C, they obtain the quoted overall SLD result.

² ABE 02G tag b and c quarks through their semileptonic decays into electrons and muons. A maximum likelihood fit is performed to extract simultaneously A_b and A_c .

³ ABE 01C tag $Z \rightarrow c\bar{c}$ events using two techniques: exclusive reconstruction of D^{*+} , D^+ and D^0 mesons and the soft pion tag for $D^{*+} \rightarrow D^0 \pi^+$. The large background from

NODE=S044AC

NODE=S044AC

NODE=S044AC

NODE=S044AC;LINKAGE=AB

NODE=S044AC;LINKAGE=G2

NODE=S044AC;LINKAGE=A

D mesons produced in $b\bar{b}$ events is separated efficiently from the signal using precision vertex information. When combining the A_c values from these two samples, care is taken to avoid double counting of events common to the two samples, and common systematic errors are properly taken into account.

A_b

This quantity is directly extracted from a measurement of the left-right forward-backward asymmetry in $b\bar{b}$ production at SLC using polarized electron beam. This double asymmetry eliminates the dependence on the Z - e - e coupling parameter A_e . OUR FIT is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the note "The Z boson" and ref. LEP-SLC 06.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.923 ± 0.020 OUR FIT				
0.9170 ± 0.0147 ± 0.0145	¹ ABE	05	SLD	$E_{cm}^{ee} = 91.24$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.907 ± 0.020 ± 0.024	48028	² ABE	03F	SLD $E_{cm}^{ee} = 91.24$ GeV
0.919 ± 0.030 ± 0.024		³ ABE	02G	SLD $E_{cm}^{ee} = 91.24$ GeV
0.855 ± 0.088 ± 0.102	7473	⁴ ABE	99L	SLD $E_{cm}^{ee} = 91.27$ GeV

¹ ABE 05 use hadronic Z decays collected during 1996–98 to obtain an enriched sample of $b\bar{b}$ events tagging on the invariant mass of reconstructed secondary decay vertices. The charge of the underlying b -quark is obtained with an algorithm that takes into account the net charge of the vertex as well as the charge of tracks emanating from the vertex and identified as kaons. This yields (25917 events) $A_b = 0.9173 \pm 0.0184 \pm 0.0173$. Taking into account all correlations with earlier results reported in ABE 03F, ABE 02G and ABE 99L, they obtain the quoted overall SLD result.

² ABE 03F obtain an enriched sample of $b\bar{b}$ events tagging on the invariant mass of a 3-dimensional topologically reconstructed secondary decay. The charge of the underlying b quark is obtained using a self-calibrating track-charge method. For the 1996–1998 data sample they measure $A_b = 0.906 \pm 0.022 \pm 0.023$. The value quoted here is obtained combining the above with the result of ABE 98i (1993–1995 data sample).

³ ABE 02G tag b and c quarks through their semileptonic decays into electrons and muons. A maximum likelihood fit is performed to extract simultaneously A_b and A_c .

⁴ ABE 99L obtain an enriched sample of $b\bar{b}$ events tagging with an inclusive vertex mass cut. For distinguishing b and \bar{b} quarks they use the charge of identified K^\pm .

TRANSVERSE SPIN CORRELATIONS IN $Z \rightarrow \tau^+ \tau^-$

The correlations between the transverse spin components of $\tau^+ \tau^-$ produced in Z decays may be expressed in terms of the vector and axial-vector couplings:

$$C_{TT} = \frac{|g_A^\tau|^2 - |g_V^\tau|^2}{|g_A^\tau|^2 + |g_V^\tau|^2}$$

$$C_{TN} = -2 \frac{|g_A^\tau| |g_V^\tau|}{|g_A^\tau|^2 + |g_V^\tau|^2} \sin(\Phi_{g_V^\tau} - \Phi_{g_A^\tau})$$

C_{TT} refers to the transverse-transverse (within the collision plane) spin correlation and C_{TN} refers to the transverse-normal (to the collision plane) spin correlation.

The longitudinal τ polarization $P_\tau (= -A_\tau)$ is given by:

$$P_\tau = -2 \frac{|g_A^\tau| |g_V^\tau|}{|g_A^\tau|^2 + |g_V^\tau|^2} \cos(\Phi_{g_V^\tau} - \Phi_{g_A^\tau})$$

Here Φ is the phase and the phase difference $\Phi_{g_V^\tau} - \Phi_{g_A^\tau}$ can be obtained using both the measurements of C_{TN} and P_τ .

C_{TT}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
1.01 ± 0.12 OUR AVERAGE				
0.87 ± 0.20 ^{+0.10} _{-0.12}	9.1k	ABREU	97G	DLPH $E_{cm}^{ee} = 91.2$ GeV
1.06 ± 0.13 ± 0.05	120k	BARATE	97D	ALEP $E_{cm}^{ee} = 91.2$ GeV

C_{TN}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.08 ± 0.13 ± 0.04	120k	¹ BARATE	97D	ALEP $E_{cm}^{ee} = 91.2$ GeV

¹ BARATE 97D combine their value of C_{TN} with the world average $P_\tau = -0.140 \pm 0.007$ to obtain $\tan(\Phi_{g_V^\tau} - \Phi_{g_A^\tau}) = -0.57 \pm 0.97$.

NODE=S044AB

NODE=S044AB

NODE=S044AB

NODE=S044AB;LINKAGE=AB

NODE=S044AB;LINKAGE=F

NODE=S044AB;LINKAGE=G2

NODE=S044AB;LINKAGE=D

NODE=S044248

NODE=S044248

NODE=S044CTT

NODE=S044CTT

NODE=S044CTN

NODE=S044CTN

NODE=S044CTN;LINKAGE=A

FORWARD-BACKWARD $e^+e^- \rightarrow f\bar{f}$ CHARGE ASYMMETRIES

These asymmetries are experimentally determined by tagging the respective lepton or quark flavor in e^+e^- interactions. Details of heavy flavor (c - or b -quark) tagging at LEP are described in the note on "The Z boson" and ref. LEP-SLC 06. The Standard Model predictions for LEP data have been (re)computed using the ZFITTER package (version 6.36) with input parameters $M_Z=91.187$ GeV, $M_{\text{top}}=174.3$ GeV, $M_{\text{Higgs}}=150$ GeV, $\alpha_s=0.119$, $\alpha^{(5)}(M_Z)=1/128.877$ and the Fermi constant $G_F=1.16637 \times 10^{-5}$ GeV $^{-2}$ (see the note on "The Z boson" for references). For non-LEP data the Standard Model predictions are as given by the authors of the respective publications.

NODE=S044280

NODE=S044280

———— $A_{FB}^{(0,e)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow e^+e^-$ ————

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note "The Z boson" and ref. LEP-SLC 06). For the Z peak, we report the pole asymmetry defined by $(3/4)A_e^2$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data.

NODE=S044Z01

NODE=S044Z01

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.45±0.25 OUR FIT				
0.89±0.44	1.57	91.2	¹ ABBIENDI	01A OPAL
1.71±0.49	1.57	91.2	ABREU	00F DLPH
1.06±0.58	1.57	91.2	ACCIARRI	00C L3
1.88±0.34	1.57	91.2	² BARATE	00C ALEP

NODE=S044Z01

¹ ABBIENDI 01A error includes approximately 0.38 due to statistics, 0.16 due to event selection systematics, and 0.18 due to the theoretical uncertainty in t -channel prediction.

NODE=S044Z01;LINKAGE=DB

² BARATE 00C error includes approximately 0.31 due to statistics, 0.06 due to experimental systematics, and 0.13 due to the theoretical uncertainty in t -channel prediction.

NODE=S044Z01;LINKAGE=A

———— $A_{FB}^{(0,\mu)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \mu^+\mu^-$ ————

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note "The Z boson" and ref. LEP-SLC 06). For the Z peak, we report the pole asymmetry defined by $(3/4)A_e A_\mu$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data.

NODE=S044Z0A

NODE=S044Z0A

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.69± 0.13 OUR FIT				
1.59± 0.23	1.57	91.2	¹ ABBIENDI	01A OPAL
1.65± 0.25	1.57	91.2	ABREU	00F DLPH
1.88± 0.33	1.57	91.2	ACCIARRI	00C L3
1.71± 0.24	1.57	91.2	² BARATE	00C ALEP

NODE=S044Z0A

• • • We do not use the following data for averages, fits, limits, etc. • • •

9 ±30	−1.3	20	³ ABREU	95M	DLPH
7 ±26	−8.3	40	³ ABREU	95M	DLPH
−11 ±33	−24.1	57	³ ABREU	95M	DLPH
−62 ±17	−44.6	69	³ ABREU	95M	DLPH
−56 ±10	−63.5	79	³ ABREU	95M	DLPH
−13 ± 5	−34.4	87.5	³ ABREU	95M	DLPH
−29.0 + 5.0 − 4.8 ±0.5	−32.1	56.9	⁴ ABE	90I	VNS
− 9.9 ± 1.5 ±0.5	−9.2	35	HEGNER	90	JADE
0.05± 0.22	0.026	91.14	⁵ ABRAMS	89D	MRK2
−43.4 ±17.0	−24.9	52.0	⁶ BACALA	89	AMY
−11.0 ±16.5	−29.4	55.0	⁶ BACALA	89	AMY
−30.0 ±12.4	−31.2	56.0	⁶ BACALA	89	AMY
−46.2 ±14.9	−33.0	57.0	⁶ BACALA	89	AMY
−29 ±13	−25.9	53.3	ADACHI	88C	TOPZ
+ 5.3 ± 5.0 ±0.5	−1.2	14.0	ADEVA	88	MRKJ
−10.4 ± 1.3 ±0.5	−8.6	34.8	ADEVA	88	MRKJ
−12.3 ± 5.3 ±0.5	−10.7	38.3	ADEVA	88	MRKJ
−15.6 ± 3.0 ±0.5	−14.9	43.8	ADEVA	88	MRKJ
− 1.0 ± 6.0	−1.2	13.9	BRAUNSCH...	88D	TASS

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=2

OCCUR=3

OCCUR=4

$-9.1 \pm 2.3 \pm 0.5$	-8.6	34.5	BRAUNSCH...	88D	TASS	OCCUR=2
$-10.6 \pm 2.2 \pm 0.5$	-8.9	35.0	BRAUNSCH...	88D	TASS	OCCUR=3
$-17.6 \pm 4.4 \pm 0.5$	-15.2	43.6	BRAUNSCH...	88D	TASS	OCCUR=4
$-4.8 \pm 6.5 \pm 1.0$	-11.5	39	BEHREND	87C	CELL	
$-18.8 \pm 4.5 \pm 1.0$	-15.5	44	BEHREND	87C	CELL	OCCUR=2
$+2.7 \pm 4.9$	-1.2	13.9	BARTEL	86C	JADE	
$-11.1 \pm 1.8 \pm 1.0$	-8.6	34.4	BARTEL	86C	JADE	OCCUR=2
$-17.3 \pm 4.8 \pm 1.0$	-13.7	41.5	BARTEL	86C	JADE	OCCUR=3
$-22.8 \pm 5.1 \pm 1.0$	-16.6	44.8	BARTEL	86C	JADE	OCCUR=4
$-6.3 \pm 0.8 \pm 0.2$	-6.3	29	ASH	85	MAC	
$-4.9 \pm 1.5 \pm 0.5$	-5.9	29	DERRICK	85	HRS	
-7.1 ± 1.7	-5.7	29	LEVI	83	MRK2	
-16.1 ± 3.2	-9.2	34.2	BRANDELIK	82C	TASS	

¹ ABBIENDI 01A error is almost entirely on account of statistics.

² BARATE 00C error is almost entirely on account of statistics.

³ ABREU 95M perform this measurement using radiative muon-pair events associated with high-energy isolated photons.

⁴ ABE 90I measurements in the range $50 \leq \sqrt{s} \leq 60.8$ GeV.

⁵ ABRAMS 89D asymmetry includes both $9 \mu^+ \mu^-$ and $15 \tau^+ \tau^-$ events.

⁶ BACALA 89 systematic error is about 5%.

NODE=S044Z0A;LINKAGE=DB

NODE=S044Z0A;LINKAGE=LB

NODE=S044Z0A;LINKAGE=H

NODE=S044Z0A;LINKAGE=AT

NODE=S044Z0A;LINKAGE=AB

NODE=S044Z0A;LINKAGE=F

———— $A_{FB}^{(0,\tau)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \tau^+\tau^-$ ————

OUR FIT is obtained using the fit procedure and correlations as determined by the LEP Electroweak Working Group (see the note “The Z boson” and ref. LEP-SLC 06). For the Z peak, we report the pole asymmetry defined by $(3/4)A_e A_\tau$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data.

NODE=S044Z0T

NODE=S044Z0T

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.88 ± 0.17 OUR FIT				
1.45 ± 0.30	1.57	91.2	¹ ABBIENDI 01A	OPAL
2.41 ± 0.37	1.57	91.2	ABREU 00F	DLPH
2.60 ± 0.47	1.57	91.2	ACCIARRI 00C	L3
1.70 ± 0.28	1.57	91.2	² BARATE 00C	ALEP

NODE=S044Z0T

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-32.8 \pm 6.4 \pm 1.5$	-32.1	56.9	³ ABE 90I	VNS	
$-8.1 \pm 2.0 \pm 0.6$	-9.2	35	HEGNER 90	JADE	
-18.4 ± 19.2	-24.9	52.0	⁴ BACALA 89	AMY	
-17.7 ± 26.1	-29.4	55.0	⁴ BACALA 89	AMY	OCCUR=2
-45.9 ± 16.6	-31.2	56.0	⁴ BACALA 89	AMY	OCCUR=3
-49.5 ± 18.0	-33.0	57.0	⁴ BACALA 89	AMY	OCCUR=4
-20 ± 14	-25.9	53.3	ADACHI 88C	TOPZ	
$-10.6 \pm 3.1 \pm 1.5$	-8.5	34.7	ADEVA 88	MRKJ	
$-8.5 \pm 6.6 \pm 1.5$	-15.4	43.8	ADEVA 88	MRKJ	OCCUR=2
$-6.0 \pm 2.5 \pm 1.0$	8.8	34.6	BARTEL 85F	JADE	
$-11.8 \pm 4.6 \pm 1.0$	14.8	43.0	BARTEL 85F	JADE	OCCUR=2
$-5.5 \pm 1.2 \pm 0.5$	-0.063	29.0	FERNANDEZ 85	MAC	
-4.2 ± 2.0	0.057	29	LEVI 83	MRK2	
-10.3 ± 5.2	-9.2	34.2	BEHREND 82	CELL	
-0.4 ± 6.6	-9.1	34.2	BRANDELIK 82C	TASS	

¹ ABBIENDI 01A error includes approximately 0.26 due to statistics and 0.14 due to event selection systematics.

² BARATE 00C error includes approximately 0.26 due to statistics and 0.11 due to experimental systematics.

³ ABE 90I measurements in the range $50 \leq \sqrt{s} \leq 60.8$ GeV.

⁴ BACALA 89 systematic error is about 5%.

NODE=S044Z0T;LINKAGE=DB

NODE=S044Z0T;LINKAGE=LB

NODE=S044Z0T;LINKAGE=AT

NODE=S044Z0T;LINKAGE=F

———— $A_{FB}^{(0,\ell)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \ell^+\ell^-$ ————

For the Z peak, we report the pole asymmetry defined by $(3/4)A_\ell^2$ as determined by the five-parameter fit to cross-section and lepton forward-backward asymmetry data assuming lepton universality. For details see the note “The Z boson” and ref. LEP-SLC 06.

NODE=S044Z0L

NODE=S044Z0L

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.71±0.10 OUR FIT				
1.45±0.17	1.57	91.2	¹ ABBIENDI	01A OPAL
1.87±0.19	1.57	91.2	ABREU	00F DLPH
1.92±0.24	1.57	91.2	ACCIARRI	00C L3
1.73±0.16	1.57	91.2	² BARATE	00C ALEP

¹ ABBIENDI 01A error includes approximately 0.15 due to statistics, 0.06 due to event selection systematics, and 0.03 due to the theoretical uncertainty in t -channel prediction.

² BARATE 00C error includes approximately 0.15 due to statistics, 0.04 due to experimental systematics, and 0.02 due to the theoretical uncertainty in t -channel prediction.

NODE=S044Z0L

NODE=S044Z0L;LINKAGE=DB

NODE=S044Z0L;LINKAGE=AC

———— $A_{FB}^{(0,u)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow u\bar{u}$ ————

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
4.0±6.7±2.8	7.2	91.2	¹ ACKERSTAFF	97T OPAL

¹ ACKERSTAFF 97T measure the forward-backward asymmetry of various fast hadrons made of light quarks. Then using SU(2) isospin symmetry and flavor independence for down and strange quarks authors solve for the different quark types.

NODE=S044Z0U

NODE=S044Z0U

NODE=S044Z0U

NODE=S044Z0U;LINKAGE=A

———— $A_{FB}^{(0,s)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow s\bar{s}$ ————

The s -quark asymmetry is derived from measurements of the forward-backward asymmetry of fast hadrons containing an s quark.

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
9.8 ±1.1 OUR AVERAGE				
10.08±1.13±0.40	10.1	91.2	¹ ABREU	00B DLPH
6.8 ±3.5 ±1.1	10.1	91.2	² ACKERSTAFF	97T OPAL

¹ ABREU 00B tag the presence of an s quark requiring a high-momentum-identified charged kaon. The s -quark pole asymmetry is extracted from the charged-kaon asymmetry taking the expected d - and u -quark asymmetries from the Standard Model and using the measured values for the c - and b -quark asymmetries.

² ACKERSTAFF 97T measure the forward-backward asymmetry of various fast hadrons made of light quarks. Then using SU(2) isospin symmetry and flavor independence for down and strange quarks authors solve for the different quark types. The value reported here corresponds then to the forward-backward asymmetry for “down-type” quarks.

NODE=S044Z0S

NODE=S044Z0S

NODE=S044Z0S

NODE=S044Z0S;LINKAGE=C

NODE=S044Z0S;LINKAGE=B

———— $A_{FB}^{(0,c)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow c\bar{c}$ ————

OUR FIT, which is obtained by a simultaneous fit to several c - and b -quark measurements as explained in the note “The Z boson” and ref. LEP-SLC 06, refers to the **Z pole** asymmetry. The experimental values, on the other hand, correspond to the measurements carried out at the respective energies.

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
7.07± 0.35 OUR FIT				
6.31± 0.93±0.65	6.35	91.26	¹ ABDALLAH	04F DLPH
5.68± 0.54±0.39	6.3	91.25	² ABBIENDI	03P OPAL
6.45± 0.57±0.37	6.10	91.21	³ HEISTER	02H ALEP
6.59± 0.94±0.35	6.2	91.235	⁴ ABREU	99Y DLPH
6.3 ± 0.9 ±0.3	6.1	91.22	⁵ BARATE	98O ALEP
6.3 ± 1.2 ±0.6	6.1	91.22	⁶ ALEXANDER	97C OPAL
8.3 ± 3.8 ±2.7	6.2	91.24	⁷ ADRIANI	92D L3

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.1 ± 3.5 ±0.5	−3.5	89.43	¹ ABDALLAH	04F DLPH
11.0 ± 2.8 ±0.7	12.3	92.99	¹ ABDALLAH	04F DLPH
− 6.8 ± 2.5 ±0.9	−3.0	89.51	² ABBIENDI	03P OPAL
14.6 ± 2.0 ±0.8	12.2	92.95	² ABBIENDI	03P OPAL
−12.4 ±15.9 ±2.0	−9.6	88.38	³ HEISTER	02H ALEP
− 2.3 ± 2.6 ±0.2	−3.8	89.38	³ HEISTER	02H ALEP
− 0.3 ± 8.3 ±0.6	0.9	90.21	³ HEISTER	02H ALEP
10.6 ± 7.7 ±0.7	9.6	92.05	³ HEISTER	02H ALEP
11.9 ± 2.1 ±0.6	12.2	92.94	³ HEISTER	02H ALEP

NODE=S044Z0C

NODE=S044Z0C

NODE=S044Z0C

OCCUR=2

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=2

OCCUR=3

OCCUR=4

OCCUR=5

OCCUR=6

12.1 ± 11.0 ± 1.0	14.2	93.90	³ HEISTER	02H	ALEP	OCCUR=7
− 4.96 ± 3.68 ± 0.53	− 3.5	89.434	⁴ ABREU	99Y	DLPH	OCCUR=2
11.80 ± 3.18 ± 0.62	12.3	92.990	⁴ ABREU	99Y	DLPH	OCCUR=3
− 1.0 ± 4.3 ± 1.0	− 3.9	89.37	⁵ BARATE	98O	ALEP	OCCUR=2
11.0 ± 3.3 ± 0.8	12.3	92.96	⁵ BARATE	98O	ALEP	OCCUR=3
3.9 ± 5.1 ± 0.9	− 3.4	89.45	⁶ ALEXANDER	97C	OPAL	
15.8 ± 4.1 ± 1.1	12.4	93.00	⁶ ALEXANDER	97C	OPAL	OCCUR=3
− 12.9 ± 7.8 ± 5.5	− 13.6	35	BEHREND	90D	CELL	
7.7 ± 13.4 ± 5.0	− 22.1	43	BEHREND	90D	CELL	OCCUR=2
− 12.8 ± 4.4 ± 4.1	− 13.6	35	ELSEN	90	JADE	
− 10.9 ± 12.9 ± 4.6	− 23.2	44	ELSEN	90	JADE	OCCUR=2
− 14.9 ± 6.7	− 13.3	35	OULD-SAADA	89	JADE	

¹ ABDALLAH 04F tag b^- and c^- quarks using semileptonic decays combined with charge flow information from the hemisphere opposite to the lepton. Enriched samples of $c\bar{c}$ and $b\bar{b}$ events are obtained using lifetime information.

NODE=S044Z0C;LINKAGE=AD

² ABBIENDI 03P tag heavy flavors using events with one or two identified leptons. This allows the simultaneous fitting of the b and c quark forward-backward asymmetries as well as the average B^0 - \bar{B}^0 mixing.

NODE=S044Z0C;LINKAGE=AB

³ HEISTER 02H measure simultaneously b and c quark forward-backward asymmetries using their semileptonic decays to tag the quark charge. The flavor separation is obtained with a discriminating multivariate analysis.

NODE=S044Z0C;LINKAGE=HH

⁴ ABREU 99Y tag $Z \rightarrow b\bar{b}$ and $Z \rightarrow c\bar{c}$ events by an exclusive reconstruction of several D meson decay modes (D^{*+} , D^0 , and D^+ with their charge-conjugate states).

NODE=S044Z0C;LINKAGE=G

⁵ BARATE 98O tag $Z \rightarrow c\bar{c}$ events requiring the presence of high-momentum reconstructed D^{*+} , D^+ , or D^0 mesons.

NODE=S044Z0C;LINKAGE=AA

⁶ ALEXANDER 97C identify the b and c events using a D/D^* tag.

NODE=S044Z0C;LINKAGE=DD

⁷ ADRIANI 92D use both electron and muon semileptonic decays.

NODE=S044Z0C;LINKAGE=A

———— $A_{FB}^{(0,b)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow b\bar{b}$ ————

NODE=S044Z0B

OUR FIT, which is obtained by a simultaneous fit to several c^- and b^- quark measurements as explained in the note “The Z boson” and ref. LEP-SLC 06, refers to the **Z pole** asymmetry. The experimental values, on the other hand, correspond to the measurements carried out at the respective energies.

NODE=S044Z0B

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
9.92 ± 0.16 OUR FIT				
9.58 ± 0.32 ± 0.14	9.68	91.231	¹ ABDALLAH	05 DLPH
10.04 ± 0.56 ± 0.25	9.69	91.26	² ABDALLAH	04F DLPH
9.72 ± 0.42 ± 0.15	9.67	91.25	³ ABBIENDI	03P OPAL
9.77 ± 0.36 ± 0.18	9.69	91.26	⁴ ABBIENDI	02I OPAL
9.52 ± 0.41 ± 0.17	9.59	91.21	⁵ HEISTER	02H ALEP
10.00 ± 0.27 ± 0.11	9.63	91.232	⁶ HEISTER	01D ALEP
7.62 ± 1.94 ± 0.85	9.64	91.235	⁷ ABREU	99Y DLPH
9.60 ± 0.66 ± 0.33	9.69	91.26	⁸ ACCIARRI	99D L3
9.31 ± 1.01 ± 0.55	9.65	91.24	⁹ ACCIARRI	98U L3
9.4 ± 2.7 ± 2.2	9.61	91.22	¹⁰ ALEXANDER	97C OPAL

NODE=S044Z0B

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.37 ± 1.43 ± 0.17	5.8	89.449	¹ ABDALLAH	05 DLPH	
10.41 ± 1.15 ± 0.24	12.1	92.990	¹ ABDALLAH	05 DLPH	OCCUR=3
6.7 ± 2.2 ± 0.2	5.7	89.43	² ABDALLAH	04F DLPH	
11.2 ± 1.8 ± 0.2	12.1	92.99	² ABDALLAH	04F DLPH	OCCUR=3
4.7 ± 1.8 ± 0.1	5.9	89.51	³ ABBIENDI	03P OPAL	OCCUR=2
10.3 ± 1.5 ± 0.2	12.0	92.95	³ ABBIENDI	03P OPAL	OCCUR=3
5.82 ± 1.53 ± 0.12	5.9	89.50	⁴ ABBIENDI	02I OPAL	
12.21 ± 1.23 ± 0.25	12.0	92.91	⁴ ABBIENDI	02I OPAL	OCCUR=3
− 13.1 ± 13.5 ± 1.0	3.2	88.38	⁵ HEISTER	02H ALEP	OCCUR=2
5.5 ± 1.9 ± 0.1	5.6	89.38	⁵ HEISTER	02H ALEP	OCCUR=3
− 0.4 ± 6.7 ± 0.8	7.5	90.21	⁵ HEISTER	02H ALEP	OCCUR=4
11.1 ± 6.4 ± 0.5	11.0	92.05	⁵ HEISTER	02H ALEP	OCCUR=5
10.4 ± 1.5 ± 0.3	12.0	92.94	⁵ HEISTER	02H ALEP	OCCUR=6
13.8 ± 9.3 ± 1.1	12.9	93.90	⁵ HEISTER	02H ALEP	OCCUR=7
4.36 ± 1.19 ± 0.11	5.8	89.472	⁶ HEISTER	01D ALEP	OCCUR=2
11.72 ± 0.97 ± 0.11	12.0	92.950	⁶ HEISTER	01D ALEP	OCCUR=3
5.67 ± 7.56 ± 1.17	5.7	89.434	⁷ ABREU	99Y DLPH	OCCUR=2
8.82 ± 6.33 ± 1.22	12.1	92.990	⁷ ABREU	99Y DLPH	OCCUR=3
6.11 ± 2.93 ± 0.43	5.9	89.50	⁸ ACCIARRI	99D L3	OCCUR=2
13.71 ± 2.40 ± 0.44	12.2	93.10	⁸ ACCIARRI	99D L3	OCCUR=3

4.95 ± 5.23 ± 0.40	5.8	89.45	⁹ ACCIARRI	98U	L3	OCCUR=2
11.37 ± 3.99 ± 0.65	12.1	92.99	⁹ ACCIARRI	98U	L3	OCCUR=3
− 8.6 ± 10.8 ± 2.9	5.8	89.45	¹⁰ ALEXANDER	97C	OPAL	
− 2.1 ± 9.0 ± 2.6	12.1	93.00	¹⁰ ALEXANDER	97C	OPAL	OCCUR=3
− 71 ± 34 ± $\frac{7}{8}$	− 58	58.3	SHIMONAKA	91	TOPZ	
− 22.2 ± 7.7 ± 3.5	− 26.0	35	BEHREND	90D	CELL	
− 49.1 ± 16.0 ± 5.0	− 39.7	43	BEHREND	90D	CELL	OCCUR=2
− 28 ± 11	− 23	35	BRAUNSCH...	90	TASS	
− 16.6 ± 7.7 ± 4.8	− 24.3	35	ELSEN	90	JADE	
− 33.6 ± 22.2 ± 5.2	− 39.9	44	ELSEN	90	JADE	OCCUR=2
3.4 ± 7.0 ± 3.5	− 16.0	29.0	BAND	89	MAC	
− 72 ± 28 ± 13	− 56	55.2	SAGAWA	89	AMY	

¹ ABDALLAH 05 obtain an enriched samples of $b\bar{b}$ events using lifetime information. The quark (or antiquark) charge is determined with a neural network using the secondary vertex charge, the jet charge and particle identification.

NODE=S044Z0B;LINKAGE=AL

² ABDALLAH 04F tag b^- and c^- quarks using semileptonic decays combined with charge flow information from the hemisphere opposite to the lepton. Enriched samples of $c\bar{c}$ and $b\bar{b}$ events are obtained using lifetime information.

NODE=S044Z0B;LINKAGE=AD

³ ABBIENDI 03P tag heavy flavors using events with one or two identified leptons. This allows the simultaneous fitting of the b and c quark forward-backward asymmetries as well as the average $B^0-\bar{B}^0$ mixing.

NODE=S044Z0B;LINKAGE=AB

⁴ ABBIENDI 02I tag $Z^0 \rightarrow b\bar{b}$ decays using a combination of secondary vertex and lepton tags. The sign of the b -quark charge is determined using an inclusive tag based on jet, vertex, and kaon charges.

NODE=S044Z0B;LINKAGE=ZQ

⁵ HEISTER 02H measure simultaneously b and c quark forward-backward asymmetries using their semileptonic decays to tag the quark charge. The flavor separation is obtained with a discriminating multivariate analysis.

NODE=S044Z0B;LINKAGE=HH

⁶ HEISTER 01D tag $Z \rightarrow b\bar{b}$ events using the impact parameters of charged tracks complemented with information from displaced vertices, event shape variables, and lepton identification. The b -quark direction and charge is determined using the hemisphere charge method along with information from fast kaon tagging and charge estimators of primary and secondary vertices. The change in the quoted value due to variation of A_{FB}^C and R_b is given as $+0.103 (A_{FB}^C - 0.0651) - 0.440 (R_b - 0.21585)$.

NODE=S044Z0B;LINKAGE=WW

⁷ ABREU 99Y tag $Z \rightarrow b\bar{b}$ and $Z \rightarrow c\bar{c}$ events by an exclusive reconstruction of several D meson decay modes (D^{*+} , D^0 , and D^+ with their charge-conjugate states).

NODE=S044Z0B;LINKAGE=XY

⁸ ACCIARRI 99D tag $Z \rightarrow b\bar{b}$ events using high p and p_T leptons. The analysis determines simultaneously a mixing parameter $\chi_b = 0.1192 \pm 0.0068 \pm 0.0051$ which is used to correct the observed asymmetry.

NODE=S044Z0B;LINKAGE=XX

⁹ ACCIARRI 98U tag $Z \rightarrow b\bar{b}$ events using lifetime and measure the jet charge using the hemisphere charge.

NODE=S044Z0B;LINKAGE=Y

¹⁰ ALEXANDER 97C identify the b and c events using a D/D^* tag.

NODE=S044Z0B;LINKAGE=FF

CHARGE ASYMMETRY IN $e^+e^- \rightarrow q\bar{q}$

NODE=S044Z0Q

Summed over five lighter flavors.

NODE=S044Z0Q

Experimental and Standard Model values are somewhat event-selection dependent. Standard Model expectations contain some assumptions on $B^0-\bar{B}^0$ mixing and on other electroweak parameters.

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
− 0.76 ± 0.12 ± 0.15		91.2	¹ ABREU	92I DLPH
4.0 ± 0.4 ± 0.63	4.0	91.3	² ACTON	92L OPAL
9.1 ± 1.4 ± 1.6	9.0	57.9	ADACHI	91 TOPZ
− 0.84 ± 0.15 ± 0.04		91	DECAMP	91B ALEP
8.3 ± 2.9 ± 1.9	8.7	56.6	STUART	90 AMY
11.4 ± 2.2 ± 2.1	8.7	57.6	ABE	89L VNS
6.0 ± 1.3	5.0	34.8	GREENSHAW	89 JADE
8.2 ± 2.9	8.5	43.6	GREENSHAW	89 JADE

NODE=S044Z0Q

¹ ABREU 92I has 0.14 systematic error due to uncertainty of quark fragmentation.

NODE=S044Z0Q;LINKAGE=B

² ACTON 92L use the weight function method on 259k selected $Z \rightarrow$ hadrons events. The systematic error includes a contribution of 0.2 due to $B^0-\bar{B}^0$ mixing effect, 0.4 due to Monte Carlo (MC) fragmentation uncertainties and 0.3 due to MC statistics. ACTON 92L derive a value of $\sin^2\theta_W^{\text{eff}}$ to be $0.2321 \pm 0.0017 \pm 0.0028$.

NODE=S044Z0Q;LINKAGE=C

OCCUR=2

CHARGE ASYMMETRY IN $p\bar{p} \rightarrow Z \rightarrow e^+e^-$

NODE=S044Z0E

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
5.2±5.9±0.4	91	ABE	91E	CDF

NODE=S044Z0E

ANOMALOUS $ZZ\gamma$, $Z\gamma\gamma$, AND ZZV COUPLINGS

A REVIEW GOES HERE – Check our WWW List of Reviews

NODE=S044270

NODE=S044270

 h_i^Y

NODE=S044ZVG

NODE=S044ZVG

Combining the LEP results properly taking into account the correlations the following 95% CL limits are derived (CERN-PH-EP/2005-051 or hep-ex/0511027):

$$\begin{aligned}
 -0.13 < h_1^Z < +0.13, & \quad -0.078 < h_2^Z < +0.071, \\
 -0.20 < h_3^Z < +0.07, & \quad -0.05 < h_4^Z < +0.12, \\
 -0.056 < h_1^\gamma < +0.055, & \quad -0.045 < h_2^\gamma < +0.025, \\
 -0.049 < h_3^\gamma < -0.008, & \quad -0.002 < h_4^\gamma < +0.034.
 \end{aligned}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=S044ZVG

• • • We do not use the following data for averages, fits, limits, etc. • • •

1	AAD	12BX ATLS	$E_{cm}^{pp} = 7$ TeV
2	ABAZOV	12S D0	$E_{cm}^{pp} = 1.96$ TeV
3	AALTONEN	11S CDF	$E_{cm}^{pp} = 1.96$ TeV
4	CHATRCHYAN 11M	CMS	$E_{cm}^{pp} = 7$ TeV
5	ABAZOV	09L D0	$E_{cm}^{pp} = 1.96$ TeV
6	ABAZOV	07M D0	$E_{cm}^{pp} = 1.96$ TeV
7	ABDALLAH	07C DLPH	$E_{cm}^{ee} = 183\text{--}208$ GeV
8	ACHARD	04H L3	$E_{cm}^{ee} = 183\text{--}208$ GeV
9	ABBIENDI,G	00C OPAL	$E_{cm}^{ee} = 189$ GeV
10	ABBOTT	98M D0	$E_{cm}^{pp} = 1.8$ TeV
11	ABREU	98K DLPH	$E_{cm}^{ee} = 161, 172$ GeV

¹ AAD 12BX study $Z\gamma$ production in pp collisions and select 142 $Z\gamma$ candidates where the Z decays to electron or muon pairs, and the photon has a transverse energy larger than 60 GeV. The expected background is 12.9 ± 4.9 events. The resulting 95% C.L. limits ranges are: $-0.028 < h_3^Z < 0.027$, $-0.022 < h_3^Z < 0.026$, $-0.00021 < h_4^Z < 0.00021$, $-0.00022 < h_4^Z < 0.00021$.

NODE=S044ZVG;LINKAGE=DA

² ABAZOV 12S study $Z\gamma$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using 6.2 fb^{-1} of data where the Z decays to electron (muon) pairs and the photon has at least 10 GeV of transverse momentum. In data, 304 (308) di-electron (di-muon) events are observed with an expected background of 255 ± 16 (285 ± 24) events. Based on the photon p_T spectrum, and including also earlier data and the $Z \rightarrow \nu\bar{\nu}$ decay mode (from ABAZOV 09L), the following 95% C.L. limits are reported: $|h_{03}^Z| < 0.026$, $|h_{04}^Z| < 0.0013$, $|h_{03}^\gamma| < 0.027$, $|h_{04}^\gamma| < 0.0014$ for a form factor scale of $\Lambda = 1.5$ TeV.

NODE=S044ZVG;LINKAGE=BA

³ AALTONEN 11S study $Z\gamma$ events in $p\bar{p}$ interactions at $\sqrt{s} = 1.96$ TeV with integrated luminosity 5.1 fb^{-1} for $Z \rightarrow e^+e^-/\mu^+\mu^-$ and 4.9 fb^{-1} for $Z \rightarrow \nu\bar{\nu}$. For the charged lepton case, the two leptons must be of the same flavor with the transverse momentum/energy of one > 20 GeV and the other > 10 GeV. The isolated photon must have $E_T > 50$ GeV. They observe 91 events with 87.2 ± 7.8 events expected from standard model processes. For the $\nu\bar{\nu}$ case they require solitary photons with $E_T > 25$ GeV and missing $E_T > 25$ GeV and observe 85 events with standard model expectation of 85.9 ± 5.6 events. Taking the form factor $\Lambda = 1.5$ TeV they derive 95% C.L. limits as $|h_3^{Z,\gamma}| < 0.022$ and $|h_4^{Z,\gamma}| < 0.0009$.

NODE=S044ZVG;LINKAGE=AL

⁴ CHATRCHYAN 11M study $Z\gamma$ production in pp collisions at $\sqrt{s} = 7$ TeV using 36 pb^{-1} pp data, where the Z decays to e^+e^- or $\mu^+\mu^-$. The total cross sections are measured for photon transverse energy $E_T^\gamma > 10$ GeV and spatial separation from charged leptons in the plane of pseudo rapidity and azimuthal angle $\Delta R(\ell, \gamma) > 0.7$ with the dilepton invariant mass requirement of $M_{\ell\ell} > 50$ GeV. The number of $e^+e^-\gamma$ and $\mu^+\mu^-\gamma$ candidates is 81 and 90 with estimated backgrounds of 20.5 ± 2.5 and 27.3 ± 3.2 events respectively. The 95% CL limits for $ZZ\gamma$ couplings are $-0.05 < h_3^Z < 0.06$ and $-0.0005 < h_4^Z < 0.0005$, and for $Z\gamma\gamma$ couplings are $-0.07 < h_3^\gamma < 0.07$ and $-0.0005 < h_4^\gamma < 0.0006$.

NODE=S044ZVG;LINKAGE=CH

⁵ ABAZOV 09L study $Z\gamma$, $Z \rightarrow \nu\bar{\nu}$ production in $p\bar{p}$ collisions at 1.96 TeV C.M. energy. They select 51 events with a photon of transverse energy E_T larger than 90 GeV, with an expected background of 17 events. Based on the photon E_T spectrum and including also Z decays to charged leptons (from ABAZOV 07M), the following 95% CL limits are reported: $|h_{30}^\gamma| < 0.033$, $|h_{40}^\gamma| < 0.0017$, $|h_{30}^Z| < 0.033$, $|h_{40}^Z| < 0.0017$.

NODE=S044ZVG;LINKAGE=AZ

⁶ ABAZOV 07M use 968 $p\bar{p} \rightarrow e^+e^-/\mu^+\mu^-\gamma X$ candidates, at 1.96 TeV center of mass energy, to tag $p\bar{p} \rightarrow Z\gamma$ events by requiring $E_T(\gamma) > 7$ GeV, lepton-gamma separation $\Delta R_{\ell\gamma} > 0.7$, and di-lepton invariant mass > 30 GeV. The cross section is in agreement with the SM prediction. Using these $Z\gamma$ events they obtain 95% C.L. limits on each h_i^V , keeping all others fixed at their SM values. They report: $-0.083 < h_{30}^Z < 0.082$, $-0.0053 < h_{40}^Z < 0.0054$, $-0.085 < h_{30}^\gamma < 0.084$, $-0.0053 < h_{40}^\gamma < 0.0054$, for the form factor scale $\Lambda = 1.2$ TeV.

NODE=S044ZVG;LINKAGE=AA

⁷ Using data collected at $\sqrt{s} = 183\text{--}208$, ABDALLAH 07C select 1,877 $e^+e^- \rightarrow Z\gamma$ events with $Z \rightarrow q\bar{q}$ or $\nu\bar{\nu}$, 171 $e^+e^- \rightarrow ZZ$ events with $Z \rightarrow q\bar{q}$ or lepton pair (except an explicit τ pair), and 74 $e^+e^- \rightarrow Z\gamma^*$ events with a $q\bar{q}\mu^+\mu^-$ or $q\bar{q}e^+e^-$ signature, to derive 95% CL limits on h_i^V . Each limit is derived with other parameters set to zero. They report: $-0.23 < h_1^Z < 0.23$, $-0.30 < h_3^Z < 0.16$, $-0.14 < h_1^\gamma < 0.14$, $-0.049 < h_3^\gamma < 0.044$.

NODE=S044ZVG;LINKAGE=AD

⁸ ACHARD 04H select 3515 $e^+e^- \rightarrow Z\gamma$ events with $Z \rightarrow q\bar{q}$ or $\nu\bar{\nu}$ at $\sqrt{s} = 189\text{--}209$ GeV to derive 95% CL limits on h_i^V . For deriving each limit the other parameters are fixed at zero. They report: $-0.153 < h_1^Z < 0.141$, $-0.087 < h_2^Z < 0.079$, $-0.220 < h_3^Z < 0.112$, $-0.068 < h_4^Z < 0.148$, $-0.057 < h_1^\gamma < 0.057$, $-0.050 < h_2^\gamma < 0.023$, $-0.059 < h_3^\gamma < 0.004$, $-0.004 < h_4^\gamma < 0.042$.

NODE=S044ZVG;LINKAGE=AC

⁹ ABBIENDI, G 00C study $e^+e^- \rightarrow Z\gamma$ events (with $Z \rightarrow q\bar{q}$ and $Z \rightarrow \nu\bar{\nu}$) at 189 GeV to obtain the central values (and 95% CL limits) of these couplings: $h_1^Z = 0.000 \pm 0.100$ ($-0.190, 0.190$), $h_2^Z = 0.000 \pm 0.068$ ($-0.128, 0.128$), $h_3^Z = -0.074^{+0.102}_{-0.103}$ ($-0.269, 0.119$), $h_4^Z = 0.046 \pm 0.068$ ($-0.084, 0.175$), $h_1^\gamma = 0.000 \pm 0.061$ ($-0.115, 0.115$), $h_2^\gamma = 0.000 \pm 0.041$ ($-0.077, 0.077$), $h_3^\gamma = -0.080^{+0.039}_{-0.041}$ ($-0.164, -0.006$), $h_4^\gamma = 0.064^{+0.033}_{-0.030}$ ($+0.007, +0.134$). The results are derived assuming that only one coupling at a time is different from zero.

NODE=S044ZVG;LINKAGE=D

¹⁰ ABBOTT 98M study $p\bar{p} \rightarrow Z\gamma + X$, with $Z \rightarrow e^+e^-$, $\mu^+\mu^-$, $\nu\bar{\nu}$ at 1.8 TeV, to obtain 95% CL limits at $\Lambda = 750$ GeV: $|h_{30}^Z| < 0.36$, $|h_{40}^Z| < 0.05$ (keeping $h_i^Z=0$), and $|h_{30}^\gamma| < 0.37$, $|h_{40}^\gamma| < 0.05$ (keeping $h_i^\gamma=0$). Limits on the CP -violating couplings are $|h_{10}^Z| < 0.36$, $|h_{20}^Z| < 0.05$ (keeping $h_i^Z=0$), and $|h_{10}^\gamma| < 0.37$, $|h_{20}^\gamma| < 0.05$ (keeping $h_i^\gamma=0$).

NODE=S044ZVG;LINKAGE=C

¹¹ ABREU 98K determine a 95% CL upper limit on $\sigma(e^+e^- \rightarrow \gamma + \text{invisible particles}) < 2.5$ pb using 161 and 172 GeV data. This is used to set 95% CL limits on $|h_{30}^\gamma| < 0.8$ and $|h_{30}^Z| < 1.3$, derived at a scale $\Lambda=1$ TeV and with $n=3$ in the form factor representation.

NODE=S044ZVG;LINKAGE=A

 f_i^γ

Combining the LEP results properly taking into account the correlations the following 95% CL limits are derived (CERN-PH-EP/2005-051 or hep-ex/0511027):

NODE=S044ZZZ

NODE=S044ZZZ

$$\begin{aligned} -0.30 < f_4^Z < +0.30, & \quad -0.34 < f_5^Z < +0.38, \\ -0.17 < f_4^\gamma < +0.19, & \quad -0.32 < f_5^\gamma < +0.36. \end{aligned}$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

NODE=S044ZZZ

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ AAD	13Z	ATLS	$E_{cm}^{pp} = 7$ TeV
² CHATRCHYAN	13B	CMS	$E_{cm}^{pp} = 7$ TeV
³ SCHAEI	09	ALEP	$E_{cm}^{ee} = 192\text{--}209$ GeV
⁴ ABAZOV	08K	D0	$E_{cm}^{p\bar{p}} = 1.96$ TeV
⁵ ABDALLAH	07C	DLPH	$E_{cm}^{ee} = 183\text{--}208$ GeV
⁶ ABBIENDI	04C	OPAL	
⁷ ACHARD	03D	L3	

- ¹ AAD 13Z study ZZ production in pp collisions at $\sqrt{s} = 7$ TeV. In the $ZZ \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-$ final state they observe a total of 66 events with an expected background of 0.9 ± 1.3 . In the $ZZ \rightarrow \ell^+ \ell^- \nu \nu$ final state they observe a total of 87 events with an expected background of 46.9 ± 5.2 . The limits on anomalous TGCs are determined using the observed and expected numbers of these ZZ events binned in p_T^Z . The 95% C.L. are as follows: for form factor scale $\Lambda = \infty$, $-0.015 < f_4^\gamma < 0.015$, $-0.013 < f_4^Z < 0.013$, $-0.016 < f_5^\gamma < 0.015$, $-0.013 < f_5^Z < 0.013$; for form factor scale $\Lambda = 3$ TeV, $-0.022 < f_4^\gamma < 0.023$, $-0.019 < f_4^Z < 0.019$, $-0.023 < f_5^\gamma < 0.023$, $-0.020 < f_5^Z < 0.019$.
- ² CHATRCHYAN 13B study ZZ production in pp collisions and select 54 ZZ candidates in the Z decay channel with electrons or muons with an expected background of 1.4 ± 0.5 events. The resulting 95% C.L. ranges are: $-0.013 < f_4^\gamma < 0.015$, $-0.011 < f_4^Z < 0.012$, $-0.014 < f_5^\gamma < 0.014$, $-0.012 < f_5^Z < 0.012$.
- ³ Using data collected in the center of mass energy range 192–209 GeV, SCHAEEL 09 select 318 $e^+ e^- \rightarrow ZZ$ events with 319.4 expected from the standard model. Using this data they derive the following 95% CL limits: $-0.321 < f_4^\gamma < 0.318$, $-0.534 < f_4^Z < 0.534$, $-0.724 < f_5^\gamma < 0.733$, $-1.194 < f_5^Z < 1.190$.
- ⁴ ABAZOV 08K search for ZZ and $Z\gamma^*$ events with $1 \text{ fb}^{-1} p\bar{p}$ data at $\sqrt{s} = 1.96$ TeV in $(ee)(ee)$, $(\mu\mu)(\mu\mu)$, $(ee)(\mu\mu)$ final states requiring the lepton pair masses to be > 30 GeV. They observe 1 event, which is consistent with an expected signal of 1.71 ± 0.15 events and a background of 0.13 ± 0.03 events. From this they derive the following limits, for a form factor (Λ) value of 1.2 TeV: $-0.28 < f_{40}^Z < 0.28$, $-0.31 < f_{50}^Z < 0.29$, $-0.26 < f_{40}^\gamma < 0.26$, $-0.30 < f_{50}^\gamma < 0.28$.
- ⁵ Using data collected at $\sqrt{s} = 183$ –208 GeV, ABDALLAH 07C select 171 $e^+ e^- \rightarrow ZZ$ events with $Z \rightarrow q\bar{q}$ or lepton pair (except an explicit τ pair), and 74 $e^+ e^- \rightarrow Z\gamma^*$ events with a $q\bar{q}\mu^+\mu^-$ or $q\bar{q}e^+e^-$ signature, to derive 95% CL limits on f_j^V . Each limit is derived with other parameters set to zero. They report: $-0.40 < f_4^Z < 0.42$, $-0.38 < f_5^Z < 0.62$, $-0.23 < f_4^\gamma < 0.25$, $-0.52 < f_5^\gamma < 0.48$.
- ⁶ ABBIENDI 04C study ZZ production in $e^+ e^-$ collisions in the C.M. energy range 190–209 GeV. They select 340 events with an expected background of 180 events. Including the ABBIENDI 00N data at 183 and 189 GeV (118 events with an expected background of 65 events) they report the following 95% CL limits: $-0.45 < f_4^Z < 0.58$, $-0.94 < f_5^Z < 0.25$, $-0.32 < f_4^\gamma < 0.33$, and $-0.71 < f_5^\gamma < 0.59$.
- ⁷ ACHARD 03D study Z -boson pair production in $e^+ e^-$ collisions in the C.M. energy range 200–209 GeV. They select 549 events with an expected background of 432 events. Including the ACCIARRI 99G and ACCIARRI 99O data (183 and 189 GeV respectively, 286 events with an expected background of 241 events) and the 192–202 GeV ACCIARRI 01I results (656 events, expected background of 512 events), they report the following 95% CL limits: $-0.48 \leq f_4^Z \leq 0.46$, $-0.36 \leq f_5^Z \leq 1.03$, $-0.28 \leq f_4^\gamma \leq 0.28$, and $-0.40 \leq f_5^\gamma \leq 0.47$.

NODE=S044ZZZ;LINKAGE=AA

NODE=S044ZZZ;LINKAGE=CH

NODE=S044ZZZ;LINKAGE=SC

NODE=S044ZZZ;LINKAGE=AB

NODE=S044ZZZ;LINKAGE=AD

NODE=S044ZZZ;LINKAGE=BI

NODE=S044ZZZ;LINKAGE=D3

ANOMALOUS W/Z QUARTIC COUPLINGS

A REVIEW GOES HERE – Check our WWW List of Reviews

a_0/Λ^2 , a_c/Λ^2

Combining published and unpublished preliminary LEP results the following 95% CL intervals for the QGCs associated with the $ZZ\gamma\gamma$ vertex are derived (CERN-PH-EP/2005-051 or hep-ex/0511027):

$$\begin{aligned} -0.008 < a_0^Z/\Lambda^2 < +0.021 \\ -0.029 < a_c^Z/\Lambda^2 < +0.039 \end{aligned}$$

VALUE	DOCUMENT ID	TECN
-------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹ ABBIENDI	04L	OPAL
² HEISTER	04A	ALEP
³ ACHARD	02G	L3

NODE=S044275

NODE=S044275

NODE=S044AQC

NODE=S044AQC

NODE=S044AQC

- ¹ ABBIENDI 04L select $20 e^+ e^- \rightarrow \nu \bar{\nu} \gamma \gamma$ acoplanar events in the energy range 180–209 GeV and $176 e^+ e^- \rightarrow q \bar{q} \gamma \gamma$ events in the energy range 130–209 GeV. These samples are used to constrain possible anomalous $W^+ W^- \gamma \gamma$ and $ZZ \gamma \gamma$ quartic couplings. Further combining with the $W^+ W^- \gamma$ sample of ABBIENDI 04B the following one-parameter 95% CL limits are obtained: $-0.007 < a_0^Z/\Lambda^2 < 0.023 \text{ GeV}^{-2}$, $-0.029 < a_c^Z/\Lambda^2 < 0.029 \text{ GeV}^{-2}$, $-0.020 < a_0^W/\Lambda^2 < 0.020 \text{ GeV}^{-2}$, $-0.052 < a_c^W/\Lambda^2 < 0.037 \text{ GeV}^{-2}$.
- ² In the CM energy range 183 to 209 GeV HEISTER 04A select $30 e^+ e^- \rightarrow \nu \bar{\nu} \gamma \gamma$ events with two acoplanar, high energy and high transverse momentum photons. The photon-photon acoplanarity is required to be $> 5^\circ$, $E_\gamma/\sqrt{s} > 0.025$ (the more energetic photon having energy $> 0.2 \sqrt{s}$), $p_{T_\gamma}/E_{\text{beam}} > 0.05$ and $|\cos \theta_\gamma| < 0.94$. A likelihood fit to the photon energy and recoil missing mass yields the following one-parameter 95% CL limits: $-0.012 < a_0^Z/\Lambda^2 < 0.019 \text{ GeV}^{-2}$, $-0.041 < a_c^Z/\Lambda^2 < 0.044 \text{ GeV}^{-2}$, $-0.060 < a_0^W/\Lambda^2 < 0.055 \text{ GeV}^{-2}$, $-0.099 < a_c^W/\Lambda^2 < 0.093 \text{ GeV}^{-2}$.
- ³ ACHARD 02G study $e^+ e^- \rightarrow Z \gamma \gamma \rightarrow q \bar{q} \gamma \gamma$ events using data at center-of-mass energies from 200 to 209 GeV. The photons are required to be isolated, each with energy $> 5 \text{ GeV}$ and $|\cos \theta| < 0.97$, and the di-jet invariant mass to be compatible with that of the Z boson (74–111 GeV). Cuts on Z velocity ($\beta < 0.73$) and on the energy of the most energetic photon reduce the backgrounds due to non-resonant production of the $q \bar{q} \gamma \gamma$ state and due to ISR respectively, yielding a total of 40 candidate events of which 8.6 are expected to be due to background. The energy spectra of the least energetic photon are fitted for all ten center-of-mass energy values from 130 GeV to 209 GeV (as obtained adding to the present analysis 130–202 GeV data of ACCIARRI 01E, for a total of 137 events with an expected background of 34.1 events) to obtain the fitted values $a_0/\Lambda^2 = 0.00_{-0.01}^{+0.02} \text{ GeV}^{-2}$ and $a_c/\Lambda^2 = 0.03_{-0.02}^{+0.01} \text{ GeV}^{-2}$, where the other parameter is kept fixed to its Standard Model value (0). A simultaneous fit to both parameters yields the 95% CL limits $-0.02 \text{ GeV}^{-2} < a_0/\Lambda^2 < 0.03 \text{ GeV}^{-2}$ and $-0.07 \text{ GeV}^{-2} < a_c/\Lambda^2 < 0.05 \text{ GeV}^{-2}$.

NODE=S044AQC;LINKAGE=AB

NODE=S044AQC;LINKAGE=HE

NODE=S044AQC;LINKAGE=C

Z REFERENCES

AAD	13Z	JHEP 1303 128	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54966
CHATRCHYAN	13B	JHEP 1301 063	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54820
AAD	12BX	PL B717 49	G. Aad <i>et al.</i>	(ATLAS Collab.)	REFID=54585
ABAZOV	12S	PR D85 052001	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=54348
CHATRCHYAN	12BN	JHEP 1212 034	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=54777
AALTONEN	11S	PRL 107 051802	T. Aaltonen <i>et al.</i>	(CDF Collab.)	REFID=16457
ABAZOV	11D	PR D84 012007	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=16465
CHATRCHYAN	11M	PL B701 535	S. Chatrchyan <i>et al.</i>	(CMS Collab.)	REFID=16645
ABAZOV	09L	PRL 102 201802	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52865
BEDDALL	09	PL B670 300	A. Beddall, A. Beddall, A. Bingul	(UGAZ)	REFID=52611
SCHAE	09	JHEP 0904 124	S. Schael <i>et al.</i>	(ALEPH Collab.)	REFID=52779
ABAZOV	08K	PRL 100 131801	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=52389
ABAZOV	07M	PL B653 378	V.M. Abazov <i>et al.</i>	(D0 Collab.)	REFID=51928
ABDALLAH	07C	EPJ C51 525	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=51839
ABDALLAH	06E	PL B639 179	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=51265
AKTAS	06	PL B632 35	A. Aktas <i>et al.</i>	(H1 Collab.)	REFID=51010
LEP-SLC	06	PRPL 427 257	ALEPH, DELPHI, L3, OPAL, SLD and working groups		REFID=51219;ERROR=1;ERROR=2
SCHAE	06A	PL B639 192	S. Schael <i>et al.</i>	(ALEPH Collab.)	REFID=51266
ABDALLAH	05	EPJ C40 1	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=50321
ABDALLAH	05C	EPJ C44 299	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=51221
ABE	05	PRL 94 091801	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=50449
ABE	05F	PR D71 112004	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=50666
ACOSTA	05M	PR D71 052002	D. Acosta <i>et al.</i>	(CDF Collab.)	REFID=50721
ABBIENDI	04B	PL B580 17	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49616
ABBIENDI	04C	EPJ C32 303	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49627
ABBIENDI	04E	PL B586 167	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49886
ABBIENDI	04G	EPJ C33 173	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49915
ABBIENDI	04L	PR D70 032005	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=50050
ABDALLAH	04F	EPJ C34 109	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49918
ABE	04C	PR D69 072003	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=49963
ACHARD	04C	PL B585 42	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49832
ACHARD	04H	PL B597 119	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=50107
HEISTER	04A	PL B602 31	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=50282
ABBIENDI	03P	PL B577 18	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=49606
ABDALLAH	03H	PL B569 129	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49548
ABDALLAH	03K	PL B576 29	J. Abdallah <i>et al.</i>	(DELPHI Collab.)	REFID=49645
ABE	03F	PRL 90 141804	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=49393
ACHARD	03D	PL B572 133	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49556
ACHARD	03G	PL B577 109	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=49808
ABBIENDI	02I	PL B546 29	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48991
ABE	02G	PRL 88 151801	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=48627
ACHARD	02G	PL B540 43	P. Achard <i>et al.</i>	(L3 Collab.)	REFID=48778
HEISTER	02B	PL B526 34	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48562
HEISTER	02C	PL B528 19	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48564
HEISTER	02H	EPJ C24 177	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48626
ABBIENDI	01A	EPJ C19 587	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48022
ABBIENDI	01G	EPJ C18 447	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48078
ABBIENDI	01K	PL B516 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48283
ABBIENDI	01N	EPJ C20 445	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48296
ABBIENDI	01O	EPJ C21 1	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=48364
ABE	01B	PRL 86 1162	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=48042
ABE	01C	PR D63 032005	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=48051
ACCIARRI	01E	PL B505 47	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=48127
ACCIARRI	01I	PL B497 23	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47915
HEISTER	01	EPJ C20 401	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48196

NODE=S044

HEISTER	01D	EPJ C22 201	A. Heister <i>et al.</i>	(ALEPH Collab.)	REFID=48508
ABBIENDI	00N	PL B476 256	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=47602
ABBIENDI,G	00C	EPJ C17 553	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=47850
ABE	00B	PRL 84 5945	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=47651
ABE	00D	PRL 85 5059	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=47880
ABREU	00	EPJ C12 225	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=47374
ABREU	00B	EPJ C14 613	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=47375
ABREU	00E	EPJ C14 585	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=47481
ABREU	00F	EPJ C16 371	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=47482
ABREU	00P	PL B475 429	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=47585
ACCIARRI	00	EPJ C13 47	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47433
ACCIARRI	00C	EPJ C16 1	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47483
ACCIARRI	00J	PL B479 79	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47620
ACCIARRI	00Q	PL B489 93	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47758
BARATE	00B	EPJ C16 597	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47376
BARATE	00C	EPJ C14 1	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47377
BARATE	00O	EPJ C16 613	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=47786
ABBIENDI	99B	EPJ C8 217	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=46595
ABBIENDI	99I	PL B447 157	G. Abbiendi <i>et al.</i>	(OPAL Collab.)	REFID=46717
ABE	99E	PR D59 052001	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=46722
ABE	99L	PRL 83 1902	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=47146
ABREU	99	EPJ C6 19	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46534
ABREU	99B	EPJ C10 415	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46594
ABREU	99J	PL B449 364	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46917
ABREU	99U	PL B462 425	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=47225
ABREU	99Y	EPJ C10 219	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=47303
ACCIARRI	99D	PL B448 152	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46733
ACCIARRI	99F	PL B453 94	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46968
ACCIARRI	99G	PL B450 281	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46991
ACCIARRI	99O	PL B465 363	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=47248
ABBOTT	98M	PR D57 R3817	B. Abbott <i>et al.</i>	(D0 Collab.)	REFID=46188
ABE	98D	PRL 80 660	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=45885
ABE	98I	PRL 81 942	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=46094
ABREU	98K	PL B423 194	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46224
ABREU	98L	EPJ C5 585	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=46265
ACCIARRI	98G	PL B431 199	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45974
ACCIARRI	98H	PL B429 387	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46064
ACCIARRI	98U	PL B439 225	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=46509
ACKERSTAFF	98A	EPJ C5 411	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46263
ACKERSTAFF	98E	EPJ C1 439	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45837
ACKERSTAFF	98O	PL B420 157	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46029
ACKERSTAFF	98Q	EPJ C4 19	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=46145
BARATE	98O	PL B434 415	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=46133
BARATE	98T	EPJ C4 557	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=46149
BARATE	98V	EPJ C5 205	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=46151
ABE	97	PRL 78 17	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=45226
ABREU	97C	ZPHY C73 243	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=45288
ABREU	97E	PL B398 207	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=45322
ABREU	97G	PL B404 194	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=45454
ACCIARRI	97D	PL B393 465	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45253
ACCIARRI	97J	PL B407 351	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45632
ACCIARRI	97L	PL B407 389	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45655
ACCIARRI	97R	PL B413 167	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=45748
ACKERSTAFF	97M	ZPHY C74 413	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45487
ACKERSTAFF	97S	PL B412 210	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45739
ACKERSTAFF	97T	ZPHY C76 387	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45785
ACKERSTAFF	97W	ZPHY C76 425	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)	REFID=45788
ALEXANDER	97C	ZPHY C73 379	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=45289
ALEXANDER	97D	ZPHY C73 569	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=45292
ALEXANDER	97E	ZPHY C73 587	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=45293
BARATE	97D	PL B405 191	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45457
BARATE	97E	PL B401 150	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45473
BARATE	97F	PL B401 163	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45474
BARATE	97H	PL B402 213	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45480
BARATE	97J	ZPHY C74 451	R. Barate <i>et al.</i>	(ALEPH Collab.)	REFID=45490
ABREU	96R	ZPHY C72 31	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44935
ABREU	96S	PL B389 405	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44969
ABREU	96U	ZPHY C73 61	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=45285
ACCIARRI	96	PL B371 126	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=44637
ADAM	96	ZPHY C69 561	W. Adam <i>et al.</i>	(DELPHI Collab.)	REFID=44627
ADAM	96B	ZPHY C70 371	W. Adam <i>et al.</i>	(DELPHI Collab.)	REFID=44629
ALEXANDER	96B	ZPHY C70 197	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=44638
ALEXANDER	96F	PL B370 185	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=44670
ALEXANDER	96N	PL B384 343	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=44901
ALEXANDER	96R	ZPHY C72 1	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=44933
BUSKULIC	96D	ZPHY C69 393	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44677
BUSKULIC	96H	ZPHY C69 379	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44703
BUSKULIC	96T	PL B384 449	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44907
BUSKULIC	96Y	PL B388 648	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44963
ABE	95J	PRL 74 2880	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=44191
ABREU	95	ZPHY C65 709 (erratum)	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44104
ABREU	95D	ZPHY C66 323	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44261
ABREU	95L	ZPHY C65 587	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44390
ABREU	95M	ZPHY C65 603	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44398
ABREU	95O	ZPHY C67 543	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44457
ABREU	95R	ZPHY C68 353	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44464
ABREU	95V	ZPHY C68 541	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44538
ABREU	95W	PL B361 207	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44542
ABREU	95X	ZPHY C69 1	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44685
ACCIARRI	95B	PL B345 589	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=44130
ACCIARRI	95C	PL B345 609	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=44132
ACCIARRI	95G	PL B353 136	M. Acciarri <i>et al.</i>	(L3 Collab.)	REFID=44281
AKERS	95C	ZPHY C65 47	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44157

AKERS	95U	ZPHY C67 389	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44380
AKERS	95W	ZPHY C67 555	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44458
AKERS	95X	ZPHY C68 1	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44459
AKERS	95Z	ZPHY C68 203	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44669
ALEXANDER	95D	PL B358 162	G. Alexander <i>et al.</i>	(OPAL Collab.)	REFID=44472
BUSKULIC	95R	ZPHY C69 15	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44813
MIYABAYASHI	95	PL B347 171	K. Miyabayashi <i>et al.</i>	(TOPAZ Collab.)	REFID=44209
ABE	94C	PRL 73 25	K. Abe <i>et al.</i>	(SLD Collab.)	REFID=43754
ABREU	94B	PL B327 386	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=43720
ABREU	94P	PL B341 109	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44067
AKERS	94P	ZPHY C63 181	R. Akers <i>et al.</i>	(OPAL Collab.)	REFID=44059
BUSKULIC	94G	ZPHY C62 179	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=43911
BUSKULIC	94J	ZPHY C62 1	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=44011
VILAIN	94	PL B320 203	P. Vilain <i>et al.</i>	(CHARM II Collab.)	REFID=43882
ABREU	93	PL B298 236	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=43224
ABREU	93I	ZPHY C59 533	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=43506
Also		ZPHY C65 709 (erratum)	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44104
ABREU	93L	PL B318 249	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=43632
ACTON	93	PL B305 407	P.D. Acton <i>et al.</i>	(OPAL Collab.)	REFID=43341
ACTON	93D	ZPHY C58 219	P.D. Acton <i>et al.</i>	(OPAL Collab.)	REFID=43378
ACTON	93E	PL B311 391	P.D. Acton <i>et al.</i>	(OPAL Collab.)	REFID=43438
ADRIANI	93	PL B301 136	O. Adriani <i>et al.</i>	(L3 Collab.)	REFID=43235
ADRIANI	93I	PL B316 427	O. Adriani <i>et al.</i>	(L3 Collab.)	REFID=43540
BUSKULIC	93L	PL B313 520	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=43524
NOVIKOV	93C	PL B298 453	V.A. Novikov, L.B. Okun, M.I. Vysotsky	(ITEP)	REFID=44105
ABREU	92I	PL B277 371	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=41986
ABREU	92M	PL B289 199	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=42151
ACTON	92B	ZPHY C53 539	D.P. Acton <i>et al.</i>	(OPAL Collab.)	REFID=41956
ACTON	92L	PL B294 436	P.D. Acton <i>et al.</i>	(OPAL Collab.)	REFID=43129
ACTON	92N	PL B295 357	P.D. Acton <i>et al.</i>	(OPAL Collab.)	REFID=43139
ADEVA	92	PL B275 209	B. Adeva <i>et al.</i>	(L3 Collab.)	REFID=41885
ADRIANI	92D	PL B292 454	O. Adriani <i>et al.</i>	(L3 Collab.)	REFID=42220
ALITTI	92B	PL B276 354	J. Alitti <i>et al.</i>	(UA2 Collab.)	REFID=41821
BUSKULIC	92D	PL B292 210	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=42210
BUSKULIC	92E	PL B294 145	D. Buskulic <i>et al.</i>	(ALEPH Collab.)	REFID=42241
DECAMP	92	PRPL 216 253	D. Decamp <i>et al.</i>	(ALEPH Collab.)	REFID=41877
ABE	91E	PRL 67 1502	F. Abe <i>et al.</i>	(CDF Collab.)	REFID=41709
ABREU	91H	ZPHY C50 185	P. Abreu <i>et al.</i>	(DELPHI Collab.)	REFID=44682
ACTON	91B	PL B273 338	D.P. Acton <i>et al.</i>	(OPAL Collab.)	REFID=41878
ADACHI	91	PL B255 613	I. Adachi <i>et al.</i>	(TOPAZ Collab.)	REFID=41667
ADEVA	91I	PL B259 199	B. Adeva <i>et al.</i>	(L3 Collab.)	REFID=44683
AKRAWY	91F	PL B257 531	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)	REFID=41638
DECAMP	91B	PL B259 377	D. Decamp <i>et al.</i>	(ALEPH Collab.)	REFID=41488
DECAMP	91J	PL B266 218	D. Decamp <i>et al.</i>	(ALEPH Collab.)	REFID=41614
JACOBSEN	91	PRL 67 3347	R.G. Jacobsen <i>et al.</i>	(Mark II Collab.)	REFID=41841
SHIMONAKA	91	PL B268 457	A. Shimonaka <i>et al.</i>	(TOPAZ Collab.)	REFID=41791
ABE	90I	ZPHY C48 13	K. Abe <i>et al.</i>	(VENUS Collab.)	REFID=41507
ABRAMS	90	PRL 64 1334	G.S. Abrams <i>et al.</i>	(Mark II Collab.)	REFID=44684
AKRAWY	90J	PL B246 285	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)	REFID=41336
BEHREND	90D	ZPHY C47 333	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=41735
BRAUNSCH...	90	ZPHY C48 433	W. Braunschweig <i>et al.</i>	(TASSO Collab.)	REFID=41736
ELSEN	90	ZPHY C46 349	E. Elsen <i>et al.</i>	(JADE Collab.)	REFID=41049
HEGNER	90	ZPHY C46 547	S. Hegner <i>et al.</i>	(JADE Collab.)	REFID=41734
STUART	90	PRL 64 983	D. Stuart <i>et al.</i>	(AMY Collab.)	REFID=41898
ABE	89	PRL 62 613	F. Abe <i>et al.</i>	(CDF Collab.)	REFID=40629
ABE	89C	PRL 63 720	F. Abe <i>et al.</i>	(CDF Collab.)	REFID=40793
ABE	89L	PL B232 425	K. Abe <i>et al.</i>	(VENUS Collab.)	REFID=41058
ABRAMS	89B	PRL 63 2173	G.S. Abrams <i>et al.</i>	(Mark II Collab.)	REFID=40962
ABRAMS	89D	PRL 63 2780	G.S. Abrams <i>et al.</i>	(Mark II Collab.)	REFID=40989
ALBAJAR	89	ZPHY C44 15	C. Albajar <i>et al.</i>	(UA1 Collab.)	REFID=40860
BACALA	89	PL B218 112	A. Bacala <i>et al.</i>	(AMY Collab.)	REFID=40805
BAND	89	PL B218 369	H.R. Band <i>et al.</i>	(MAC Collab.)	REFID=40807
GREENSHAW	89	ZPHY C42 1	T. Greenshaw <i>et al.</i>	(JADE Collab.)	REFID=40853
OULD-SAADA	89	ZPHY C44 567	F. Ould-Saada <i>et al.</i>	(JADE Collab.)	REFID=40965
SAGAWA	89	PRL 63 2341	H. Sagawa <i>et al.</i>	(AMY Collab.)	REFID=40979
ADACHI	88C	PL B208 319	I. Adachi <i>et al.</i>	(TOPAZ Collab.)	REFID=40766
ADEVA	88	PR D38 2665	B. Adeva <i>et al.</i>	(Mark-J Collab.)	REFID=40686
BRAUNSCH...	88D	ZPHY C40 163	W. Braunschweig <i>et al.</i>	(TASSO Collab.)	REFID=40697
ANSARI	87	PL B186 440	R. Ansari <i>et al.</i>	(UA2 Collab.)	REFID=40199
BEHREND	87C	PL B191 209	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=40203
BARTEL	86C	ZPHY C30 371	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=40179
Also		ZPHY C26 507	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=10047
Also		PL 108B 140	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=10033
ASH	85	PRL 55 1831	W.W. Ash <i>et al.</i>	(MAC Collab.)	REFID=10046
BARTEL	85F	PL 161B 188	W. Bartel <i>et al.</i>	(JADE Collab.)	REFID=10342
DERRICK	85	PR D31 2352	M. Derrick <i>et al.</i>	(HRS Collab.)	REFID=10049
FERNANDEZ	85	PRL 54 1624	E. Fernandez <i>et al.</i>	(MAC Collab.)	REFID=10347
LEVI	83	PRL 51 1941	M.E. Levi <i>et al.</i>	(Mark II Collab.)	REFID=41536
BEHREND	82	PL 114B 282	H.J. Behrend <i>et al.</i>	(CELLO Collab.)	REFID=10324
BRANDELIK	82C	PL 110B 173	R. Brandelik <i>et al.</i>	(TASSO Collab.)	REFID=41537